

DAFTAR PUSTAKA

- Acar, Y., Doğan, L., Güven, H. E., Aksel, B., Karaman, N., Özaslan, C., & Gülçelik, M. A. (2017). Bethesda Made It Clearer: A Review of 542 Patients in a Single Institution. *Oncology Research and Treatment*, 40(5), 277–280. <https://doi.org/10.1159/000460298>
- Aromataris, E., & Munn, Z. (Eds.). (2020). *JBI Manual for Evidence Synthesis* (Issue April). JBI. <https://doi.org/10.46658/JBIMES-20-01>
- Arul, P., & Masilamani, S. (2015). A correlative study of solitary thyroid nodules using the bethesda system for reporting thyroid cytopathology. *Journal of Cancer Research and Therapeutics*, 11(3), 617–622. <https://doi.org/10.4103/0973-1482.157302>
- Ashorobi, D., & Lopez, P. P. (2021). *Follicular Thyroid Cancer*. StatPearls.
- Aydoğan, B. İ., Şahin, M., Ceyhan, K., Deniz, O., Demir, Ö., Emral, R., Tonyukuk Gedik, V., Uysal, A. R., & Çorapçıoğlu, D. (2019). The influence of thyroid nodule size on the diagnostic efficacy and accuracy of ultrasound guided fine-needle aspiration cytology. *Diagnostic Cytopathology*, 47(7), 682–687. <https://doi.org/10.1002/dc.24170>
- Ayub, A. D. (2017). Korelasi Aspirasi Jarum Halus Dengan Histopatologi Akhir Pada Neoplasma Tiroid Di Bagian Histopatologi Anatomi Rumah Sakit Universitas Hasanuddin Makassar Periode 2015-2016. *Universitas Hasanuddin Makassar*. http://digilib.unhas.ac.id/uploaded_files/temporary/DigitalCollection/ZTY3ZWJINWQ4ZTZINDE0NmUwZTA0ZTFjYTNIZTQ4YThiNDk0MDJhOA==.pdf
- Bahaj, A. S., Alkaff, H. H., Melebari, B. N., Melebari, A. N., Sayed, S. I., Mujtaba, S. S., Abdelmonim, S. K., Alghamdi, S. A., Alghamdi, F. E., Abu Suliman, O. A., Alessa, M. A., Abi Sheffah, F. R., Al-Tammas, A. H., Al-Zahrani, R. A., Marglani, O. A., Heaphy, J. C., Bawazir, O. A., & Alherabi, A. Z. (2020). Role of fine-needle aspiration cytology in evaluating thyroid nodules A retrospective study from a tertiary care center of Western Region, Saudi Arabia. *Saudi Medical Journal*, 41(10), 1098–1103. <https://doi.org/10.15537/SMJ.2020.10.25417>
- Bailey, S., & Wallwork, B. (2018). Differentiating between benign and malignant thyroid nodules: An evidence-based approach in general practice. *Australian Journal of General Practice*, 47(11), 770–774. <https://doi.org/10.31128/ajgp-03-18-4518>
- Bestepe, N., Ozdemir, D., Tam, A. A., Dellal, F. D., Kilicarslan, A., Parlak, O.,

- Ersoy, R., & Cakir, B. (2016). Malignancy risk and false-negative rate of fine needle aspiration cytology in thyroid nodules ≥ 4.0 cm. *Surgery (United States)*, 160(2), 405–412. <https://doi.org/10.1016/j.surg.2016.03.019>
- Bohacek, L., Milas, M., Mitchell, J., Siperstein, A., & Berber, E. (2012). Diagnostic accuracy of surgeon-performed ultrasound-guided fine-needle aspiration of thyroid nodules. *Annals of Surgical Oncology*, 19(1), 45–51. <https://doi.org/10.1245/s10434-011-1807-z>
- Bresler, A., Mehta, V., Schiff, B. A., Smith, R. V., Khader, S., Ramos-Rivera, G., Lin, J., Libutti, S. K., Laird, A. M., & Ow, T. J. (2019). Comparison of Bethesda cytopathology classification to surgical pathology across racial-ethnic groups. *Head and Neck*, 41(7), 2340–2345. <https://doi.org/10.1002/hed.25707>
- Burman, K. D., & Wartofsky, L. (2015). CLINICAL PRACTICE. Thyroid Nodules. *The New England Journal of Medicine*, 373(24), 2347–2356. <https://doi.org/10.1056/NEJMcp1415786>
- Campbell, J. M., Klugar, M., Ding, S., Carmody, D. P., Hakonen, S. J., Jadotte, Y. T., White, S., & Munn, Z. (2015). Diagnostic test accuracy: Methods for systematic review and meta-analysis. *International Journal of Evidence-Based Healthcare*, 13(3), 154–162. <https://doi.org/10.1097/XEB.0000000000000061>
- Chang, H., Lee, E., Lee, H., Choi, J., Kim, A., & Kim, B. hui. (2013). Comparison of diagnostic values of thyroid aspiration samples using liquid-based preparation and conventional smear: One-year experience in a single institution. *Apmis*, 121(2), 139–145. <https://doi.org/10.1111/j.1600-0463.2012.02944.x>
- Cibas, E. S., & Ali, S. Z. (2009). The Bethesda system for reporting thyroid cytopathology. *American Journal of Clinical Pathology*, 132(5), 658–665. <https://doi.org/10.1309/AJCPHLWMI3JV4LA>
- Cibas, E. S., Baloch, Z. W., Fellegara, G., LiVolsi, V. A., Raab, S. S., Rosai, J., Diggans, J., Friedman, L., Kennedy, G. C., Kloos, R. T., Lanman, R. B., Mandel, S. J., Sindy, N., Steward, D. L., Zeiger, M. A., Haugen, B. R., & Alexander, E. K. (2013). A Prospective Assessment Defining the Limitations of Thyroid Nodule Pathologic Evaluation. *Annals of Internal Medicine*, 159(5), 325. <https://doi.org/10.7326/0003-4819-159-5-201309030-00006>
- de Cristo, A. P., Goldstein, H. F., Faccin, C. S., Maia, A. L., & Graudenz, M. S. (2016). Increasing diagnostic effectiveness of thyroid nodule evaluation by implementation of cell block preparation in routine US-FNA analysis. *Archives of Endocrinology and Metabolism*, 60(4), 367–373. <https://doi.org/10.1590/2359-3997000000180>

- DeVita, V. T., Lawrence, T. S., & Rosenberg, S. A. (2019). *Devita, Hellman, and Rosenberg's cancer: principles & practice of oncology* (11th ed.). Wolters Kluwer.
- Dong, Y. J., Mao, M. J., Zhan, W. W., Zhou, J. Q., Zhou, W., Yao, J. J., Hu, Y. Y., Wang, Y., & Ye, T. J. (2018). Size and ultrasound features affecting results of ultrasound-guided fine-needle aspiration of thyroid nodules. *Journal of Ultrasound in Medicine*, 37(6), 1367–1377. <https://doi.org/10.1002/jum.14472>
- Drake, R. L., Vogl, A. W., & Mitchell, A. W. M. (2020). *Gray's anatomy for students* (4th ed.). Elsevier.
- Erkinuresin, T., & Demirci, H. (2019). Diagnostic accuracy of fine Needle Aspiration Cytology in thyroid nodules. *Diagnosis*, 7(1), 61–66. <https://doi.org/10.1515/dx-2019-0039>
- Eusebi, P. (2013). Diagnostic accuracy measures. *Cerebrovascular Diseases*, 36(4), 267–272. <https://doi.org/10.1159/000353863>
- Evranos, B., Polat, S. B., Baser, H., Ozdemir, D., Kilicarslan, A., Yalcin, A., Ersoy, R., & Cakir, B. (2017). Bethesda classification is a valuable guide for fine needle aspiration reports and highly predictive especially for diagnosing aggressive variants of papillary thyroid carcinoma. *Cytopathology*, 28(4), 259–267. <https://doi.org/10.1111/cyt.12384>
- Fernandes, V. T., Magarey, M. J. R., Kamdar, D. P., & Freeman, J. L. (2016). Surgeon performed ultrasound-guided fine-needle aspirates of the thyroid: 1067 biopsies and learning curve in a teaching center. *Head & Neck*, 38(S1), E1281–E1284. <https://doi.org/10.1002/hed.24212>
- Gao, L., Ma, B., Zhou, L., Wang, Y., Yang, S., Qu, N., Gao, Y., & Ji, Q. (2017). The impact of presence of Hashimoto's thyroiditis on diagnostic accuracy of ultrasound-guided fine-needle aspiration biopsy in subcentimeter thyroid nodules: A retrospective study from FUSCC. *Cancer Medicine*, 6(5), 1014–1022. <https://doi.org/10.1002/cam4.997>
- Globocan Observatory, W. (2019). Cancer Today - World. *International Agency for Research on Cancer*, 876, 2018–2019. <https://gco.iarc.fr/today/data/factsheets/populations/900-world-factsheets.pdf>
- Godazandeh, G., Kashi, Z., Zargarnataj, S., Fazli, M., Ebadi, R., & Kerdabadi, E. H. (2016). Evaluation the relationship between thyroid nodule size with malignancy and accuracy of fine needle aspiration biopsy (FNAB). *Acta Informatica Medica*, 24(5), 347–351. <https://doi.org/10.5455/aim.2016.24.347-350>

- Grani, G., Sponziello, M., Pecce, V., Ramundo, V., & Durante, C. (2020). Contemporary Thyroid Nodule Evaluation and Management. *The Journal of Clinical Endocrinology & Metabolism*, 105(9), 2869–2883. <https://doi.org/10.1210/clinem/dgaa322>
- Guo, H. Q., Zhang, Z. H., Zhao, H., Niu, L. J., Chang, Q., & Pan, Q. J. (2015). Factors influencing the reliability of thyroid fine-needle aspiration: Analysis of thyroid nodule size, guidance mode for aspiration and preparation method. *Acta Cytologica*, 59(2), 169–174. <https://doi.org/10.1159/000381412>
- Hall, J. E., & Guyton, A. C. (2016). *Guyton and Hall textbook of medical physiology* (13th ed.). Elsevier.
- Heilo, A., Sigstad, E., & Grøholt, K. (2010). *Atlas of Thyroid Lesions* (2011th ed.). Springer. <https://doi.org/10.1007/978-1-4419-6010-8>
- Iqbal, J., Aziz, O. B. A., Ahmad, N., Tariq, M., & Anwar, Z. (2016). Diagnostic accuracy of fine needle aspiration cytology in the diagnosis of malignant solitary thyroid nodule. *Pakistan Armed Forces Medical Journal*, 66(4). <https://www.pafmj.org/index.php/PAFMJ/article/view/637>
- Jiang, K., Zhou, J., Lei, J., Liu, Y., Li, J., Su, X., Li, Z., Wei, T., Jiang, Y., & Zhu, J. (2018). Cell block is a valuable adjunct to conventional smear for thyroid fine needle aspiration: 2395 cases with histological correlation. *Cytopathology*, 29(6), 525–530. <https://doi.org/10.1111/cyt.12633>
- Jinjh, M., Faisal, F., Abdalla, K., Majeed, M., Achakzai, A. A., Heffron, C., McCarthy, J., & Redmond, H. P. (2020). Association between thyroid nodule size and malignancy rate. *Annals of the Royal College of Surgeons of England*, 102(1), 43–48. <https://doi.org/10.1308/rcsann.2019.0156>
- K, S., T M, N., D, M., Mahajan, R., Kumar, P., George, G. K., & Boban, M. (2015). a Study of Correlation of Preoperative Fine Needle Aspiration Cytology (Fnac) With Histopathological Examination (Hpe) in Goitre. *Journal of Evolution of Medical and Dental Sciences*, 4(89), 15414–15417. <https://doi.org/10.14260/jemds/2015/2198>
- Kaliszewski, K., Diakowska, D., Wojtczak, B., Strutyńska-Karpińska, M., Domosławski, P., Sutkowski, K., Głód, M., Balcerzak, W., Forkasiewicz, Z., & Łukieńczuk, T. (2016). Fine-needle aspiration biopsy as a preoperative procedure in patients with malignancy in solitary and multiple thyroid nodules. *PLoS ONE*, 11(1), 1–13. <https://doi.org/10.1371/journal.pone.0146883>
- Karadeniz, E., Yur, M., Temiz, A., & Akçay, M. N. (2019). Malignancy risk for thyroid nodules larger than 4 cm and diagnostic reliability of ultrasound-guided FNAB results. *Turkish Journal of Surgery*, 35(1), 013–018. <https://doi.org/10.5578/turkjsurg.4029>

- Kim, S. Y., Lee, H. S., Moon, J., Kim, E. K., Moon, H. J., Yoon, J. H., & Kwak, J. Y. (2017). Fine-needle aspiration versus core needle biopsy for diagnosis of thyroid malignancy and neoplasm: a matched cohort study. *European Radiology*, 27(2), 801–811. <https://doi.org/10.1007/s00330-016-4424-1>
- Lale, S. A., Morgenstern, N. N., Chiara, S., & Wasserman, P. (2015). Fine needle aspiration of thyroid nodules in the pediatric population: A 12-year cyto-histological correlation experience at north shore-long island jewish health system. *Diagnostic Cytopathology*, 43(8), 598–604. <https://doi.org/10.1002/dc.23265>
- Lee, Y.-B., Cho, Y. Y., Jang, J. Y., Kim, T. H., Jang, H. W., Chung, J. H., Oh, Y. L., & Kim, S. W. (2017). Current status and diagnostic values of the Bethesda system for reporting thyroid cytopathology in a papillary thyroid carcinoma-prevalent area. *Head & Neck*, 39(2), 269–274. <https://doi.org/10.1002/hed.24578>
- Limaiem, F., Rehman, A., & Mazzoni, T. (2021). *Papillary Thyroid Carcinoma*. StatPearls.
- Lloyd, R. V., Osamura, R. Y., Klöppel, G., & Rosai, J. (2017). *World Health Organization Classification of Tumours* (4th ed.). International Agency for Research on Cancer.
- Lodewijk, L., Vriens, M. R., Vorselaars, W. M. C. M., van der Meij, N. T. M., Kist, J. W., Barentsz, M. W., Verkooijen, H. M., Rinkes, I. H. M. B., & Valk, G. D. (2016). Same-Day Fine-Needle Aspiration Cytology Diagnosis for Thyroid Nodules Achieves Rapid Anxiety Decrease and High Diagnostic Accuracy. *Endocrine Practice : Official Journal of the American College of Endocrinology and the American Association of Clinical Endocrinologists*, 22(5), 561–566. <https://doi.org/10.4158/EP151036.OR>
- Machała, E., Sopiński, J., Iavorska, I., & Kołomecki, K. (2018). Correlation of Fine Needle Aspiration Cytology of Thyroid Gland with Histopathological Results. *Polish Journal of Surgery*, 90(5), 1–5. <https://doi.org/10.5604/01.3001.0012.4712>
- Magister, M. J., Chaikhoutdinov, I., Schaefer, E., Williams, N., Saunders, B., & Goldenberg, D. (2015). Association of thyroid nodule size and Bethesda class with rate of malignant disease. *JAMA Otolaryngology - Head and Neck Surgery*, 141(12), 1089–1095. <https://doi.org/10.1001/jamaoto.2015.1451>
- Master, S. R., & Burns, B. (2021). *Medullary Thyroid Cancer*. StatPearls.
- Mayocean, N., Waters, P. S., Kaim Khani, T. Y., Kerin, M. J., & Quill, D. (2016). FNAC and frozen section correlations with definitive histology in thyroid diseases. *European Archives of Oto-Rhino-Laryngology*, 273(8), 2181–2184.

<https://doi.org/10.1007/s00405-015-3742-2>

- Mescher, A. L. (2013). *Junqueiras Basic Histology Text and Atlas* (13th ed.). McGraw-Hill Education.
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., Shekelle, P., & Stewart, L. A. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews*, 4(1), 1. <https://doi.org/10.1186/2046-4053-4-1>
- Nagarajan, N., Schneider, E. B., Ali, S. Z., Zeiger, M. A., & Olson, M. T. (2015). How do liquid-based preparations of thyroid fine-needle aspiration compare with conventional smears? An analysis of 5475 specimens. *Thyroid*, 25(3), 308–313. <https://doi.org/10.1089/thy.2014.0394>
- Norlén, O., Charlton, A., Sarkis, L. M., Henwood, T., Shun, A., Gill, A. J., & Delbridge, L. (2015). Risk of malignancy for each Bethesda class in pediatric thyroid nodules. *Journal of Pediatric Surgery*, 50(7), 1147–1149. <https://doi.org/10.1016/j.jpedsurg.2014.10.046>
- Norris, J. J., & Farni, F. (2021). *Follicular Adenoma*. StatPearls.
- Nugraha, M. T. (2020). Perbandingan Akurasi Nilai Fine Needle Aspiration Biopsy (FNAB) Dengan Biopsi Patologi Anatomi Dalam Diagnostik Nodul Tiroid di RSUD Pare Kediri Tahun 2015-2019. *Universitas Muhammadiyah Surabaya*.
- Orell, S. R., & Sterrett, G. F. (2012). *Orell and sterret's fine needle aspiration cytology* (5th ed.). Elsevier.
- Ozdemir, D., Bestepe, N., Faki, S., Kilicarslan, A., Parlak, O., Ersoy, R., & Cakir, B. (2017). Comparison of thyroid fine needle aspiration biopsy results before and after implementation of Bethesda classification. *Cytopathology*, 28(5), 400–406. <https://doi.org/10.1111/cyt.12437>
- Ozluk, Y., Pehlivan, E., Gulluoglu, M. G., Poyanli, A., Salmaslioglu, A., Colak, N., Kapran, Y., & Yilmazbayhan, D. (2011). The use of the Bethesda terminology in thyroid fine-needle aspiration results in a lower rate of surgery for nonmalignant nodules: A report from a reference center in Turkey. *International Journal of Surgical Pathology*, 19(6), 761–771. <https://doi.org/10.1177/1066896911415667>
- Prades, J. M., Querat, C., Dumollard, J. M., Richard, C., Gavid, M., Timoshenko, A. A., Peoc'h, M., & Martin, C. (2013). Thyroid nodule surgery: Predictive diagnostic value of fine-needle aspiration cytology and frozen section. *European Annals of Otorhinolaryngology, Head and Neck Diseases*, 130(4), 195–199. <https://doi.org/10.1016/j.anrol.2012.12.005>
- Prasetyo, Z. A. (2012). Uji Diagnostik Fnab (Fine Needle Aspiration Biopsy)

- Dibandingkan Dengan Biopsi Patologi Anatomi Dalam Mendiagnosis Karsinoma Tiroid. *Jurnal Kedokteran Diponegoro*.
- Putri, D. R. M. S. (2018). Nilai Diagnostik Biopsi Aspirasi Jarum Halus Untuk Membedakan Tumor Jinak Dan Tumor Ganas Tiroid Dengan Baku Emas Histopatologi. *Universitas Brawijaya Malang*.
- Raguin, T., Schneegans, O., Rodier, J.-F., Volkmar, P.-P., Sauleau, E., Debry, C., Debonnecaze, G., Ghnassia, J.-P., & Dupret-Bories, A. (2017). Value of fine-needle aspiration in evaluating large thyroid nodules. *Head & Neck*, 39(1), 32–36. <https://doi.org/10.1002/hed.24524>
- Regita, A. N. (2018). *NILAI DIAGNOSTIK FINE NEEDLE ASPIRATION BIOPSY (FNAB) UNTUK MEMBEDAKAN TUMOR JINAK DAN TUMOR GANAS TULANG DENGAN BAKU EMAS HISTOPATOLOGI*. Universitas Brawijaya Malang.
- Sarkis, L. M., Norlen, O., Aniss, A., Watson, N., Delbridge, L. W., Sidhu, S. B., Sywak, M. S., & Gill, A. J. (2014). The Australian experience with the Bethesda classification system for thyroid fine needle aspiration biopsies. *Pathology*, 46(7), 592–595. <https://doi.org/10.1097/PAT.0000000000000168>
- Sharma, R., Verma, N., Kaushal, V., Sharma, D., & Sharma, D. (2017). Diagnostic accuracy of fine-needle aspiration cytology of thyroid gland lesions: A study of 200 cases in Himalayan belt. *Journal of Cancer Research and Therapeutics*, 13(3), 451–455. <https://doi.org/10.4103/0973-1482.164702>
- Sherwood, L. (2016). *Human physiology. from cells to systems* (9th ed.). Cengage Learning.
- Shin, I., Kim, E. K., Moon, H. J., Yoon, J. H., Park, V. Y., Lee, S. E., Lee, H. S., & Kwak, J. Y. (2020). Core-needle biopsy does not show superior diagnostic performance to fine-needle aspiration for diagnosing thyroid nodules. *Yonsei Medical Journal*, 61(2), 161–168. <https://doi.org/10.3349/ymj.2020.61.2.161>
- Singh, A., Pant, S., Jain, R., & Kumar, S. (2017). A Prospective Study on Assessing the Reliability of Fine Needle Aspiration Cytology in Outcomes of Thyroid Lesions. *International Archives of BioMedical and Clinical Research*, 3(3), 33–35. <https://doi.org/10.21276/iabcr.2017.3.3.9>
- Siswanto, S. (2012). Systematic Review as a Research Method to Synthesize Research Results (Introduction). *Buletin Penelitian Sistem Kesehatan*, 13(4).
- Sunder, K. S., & Khan, M. I. (2017). Role of FNAC in diagnosis of Thyroid lesions. *Journal of Contemporary Medicine and Dentistry*, 5(1), 30–34.
- Sung, J. Y., Na, D. G., Kim, K. S., Yoo, H., Lee, H., Kim, J. H., & Baek, J. H. (2012). Diagnostic accuracy of fine-needle aspiration versus core-needle

- biopsy for the diagnosis of thyroid malignancy in a clinical cohort. *European Radiology*, 22(7), 1564–1572. <https://doi.org/10.1007/s00330-012-2405-6>
- Taha, I., Al-Thani, H., El-Menyar, A., Asim, M., Al-Sulaiti, M., & Tabeb, A. (2020). Diagnostic accuracy of preoperative palpation- versus ultrasound-guided thyroid fine needle aspiration cytology: an observational study. *Postgraduate Medicine*, 132(5), 465–472. <https://doi.org/10.1080/00325481.2020.1741298>
- Thewjitcharoen, Y., Butadej, S., Nakasatien, S., Chotwanvirat, P., Porramatikul, S., Krittiyawong, S., Lekpittaya, N., & Himathongkam, T. (2019). Incidence and malignancy rates classified by The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC) – An 8-year tertiary center experience in Thailand. *Journal of Clinical and Translational Endocrinology*, 16(November 2018), 100175. <https://doi.org/10.1016/j.jcte.2018.12.004>
- Ugurluoglu, C., Dobur, F., Karabagli, P., & Celik, Z. E. (2015). Fine needle aspiration biopsy of thyroid nodules: Cytologic and histopathologic correlation of 1096 patients. *International Journal of Clinical and Experimental Pathology*, 8(11), 14800–14805.
- Wharry, L. I., McCoy, K. L., Stang, M. T., Armstrong, M. J., Lebeau, S. O., Tublin, M. E., Sholosh, B., Silbermann, A., Ohori, N. P., Nikiforov, Y. E., Hodak, S. P., Carty, S. E., & Yip, L. (2014). Thyroid nodules (≥ 4 cm): Can ultrasound and cytology reliably exclude cancer? *World Journal of Surgery*, 38(3), 614–621. <https://doi.org/10.1007/s00268-013-2261-9>
- Wong, R., Farrell, S. G., & Grossmann, M. (2018). Thyroid nodules: Diagnosis and management. *Medical Journal of Australia*, 209(2), 92–98. <https://doi.org/10.5694/mja17.01204>
- World Health Organization. (2019). Indonesia Source GLOBOCAN 2018. *International Agency for Research on Cancer*, 256, 1–2. <http://gco.iarc.fr/>