

## DAFTAR PUSTAKA

- Abd El Latif, A. *et al.* (2014) 'Moringa oleifera leaf extract ameliorates alloxan-induced diabetes in rats by regeneration of  $\beta$  cells and reduction of pyruvate carboxylase expression', *Biochemistry and Cell Biology*, 92(5). doi: 10.1139/bcb-2014-0081.
- Abd Eldaim, M. A., Shaban Abd Elrasoul, A. and Abd Elaziz, S. A. (2017) 'An aqueous extract from Moringa oleifera leaves ameliorates hepatotoxicity in alloxan-induced diabetic rats', *Biochemistry and Cell Biology*, 95(4). doi: 10.1139/bcb-2016-0256.
- Adejoh, I. P. *et al.* (2015) 'Assessment of biochemical and histopathological changes in diabetic wistar rats co-administered ethanolic leaf extract of Moringa oleifera and metformin', *International Journal of Biological & Pharmaceutical Research*, 6(12).
- Adisakwattana, S. and Chanathong, B. (2011) ' $\alpha$ -glucosidase inhibitory activity and lipid-lowering mechanisms of Moringa oleifera leaf extract', *European Review for Medical and Pharmacological Sciences*, 15(7).
- Aida, K. *et al.* (2018) 'Distinct Inflammatory Changes of the Pancreas of Slowly Progressive Insulin-dependent (Type 1) Diabetes', *Pancreas*, 47(9), pp. 1101–1109. doi: 10.1097/MPA.0000000000001144.
- Aisyah, L. S. *et al.* (2017) 'Flavonoid compounds from the leaves of Kalanchoe prolifera and their cytotoxic activity against P-388 murine leukemia cells', *Natural Product Sciences*, 23(2). doi: 10.20307/nps.2017.23.2.139.
- Al-Malki, A. L. and El Rabey, H. A. (2015) 'The antidiabetic effect of low doses of moringa oleifera lam. Seeds on streptozotocin induced diabetes and diabetic nephropathy in male rats', *BioMed Research International*, 2015. doi: 10.1155/2015/381040.
- Alberti, K. G. M. M. and Zimmet, P. Z. (1998) 'Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus. Provisional report of a WHO consultation', *Diabetic Medicine*, 15(7). doi: 10.1002/(SICI)1096-9136(199807)15:7<539::AID-DIA668>3.0.CO;2-S.
- Alethea, T. and Ramadhian, M. R. (2015) 'Efek Antidiabetik pada Daun Kelor', *Jurnal Majority*, Vol 4(No 9).
- Ali, F. T., Hassan, N. S. and Abd Rabou, R. R. (2015) 'Potential activity of Moringa Oleifera leaf extract and some active ingredients against diabetes in rats', *International Journal of Scientific & Engineering Research*, 6(5).

- Aminah, S., Ramdhan, T. and Yanis, M. (2015) ‘Syarifah Am inah et. al. : Kandungan Nutrisi dan Sifat Fungsional Tanaman Kelor (Moringa oleifera)’, *Buletin Pertanian Perkotaan*, 5(30).
- Amzu, E. (2015) ‘Kampung Konservasi Kelor: Upaya Mendukung Gerakan Nasional Sadar Gizi Dan Mengatasi Malnutrisi Di Indonesia’, *Risalah Kebijakan Pertanian Dan Lingkungan: Rumusan Kajian Strategis Bidang Pertanian dan Lingkungan*, 1(2). doi: 10.20957/jkebijakan.v1i2.10298.
- Auliya, D., Saptadi, D. and Kuswanto, K. (2019) ‘Eksplorasi Tanaman Kelor (Moringa oleifera Lam.) di Kabupaten Banyuwangi Jawa Timur’, *Jurnal Produksi Tanaman*, 6(11). doi: 10.21176/PROTAN.V6I11.1012.
- Bahadoran, Z., Mirmiran, P. and Azizi, F. (2013) ‘Dietary polyphenols as potential nutraceuticals in management of diabetes: A review’, *Journal of Diabetes and Metabolic Disorders*. doi: 10.1186/2251-6581-12-43.
- Bamagous, G. A. et al. (2018) ‘Antidiabetic and antioxidant activity of ethyl acetate extract fraction of Moringa oleifera leaves in streptozotocin-induced diabetes rats via inhibition of inflammatory mediators’, *Asian Pacific Journal of Tropical Biomedicine*, 8(6). doi: 10.4103/2221-1691.235327.
- Baset, M. et al. (2020) ‘Anti-diabetic effects of fenugreek (Trigonella foenum-graecum): A comparison between oral and intraperitoneal administration - an animal study’, *International Journal of Functional Nutrition*. doi: 10.3892/ijfn.2020.2.
- Bekoe, E. O., Jibira, Y. and Agyei, G. A. (2020) ‘Physicochemical, phytochemical, heavy metal and microbiological analysis of moringa oleifera lam. leaves’, *Pharmacognosy Journal*, 12(6). doi: 10.5530/pj.2020.12.211.
- Bhattacharya, A. et al. (2018) ‘A Review of the Phytochemical and Pharmacological Characteristics of Moringa oleifera’, *Journal of Pharmacy & Bioallied Sciences*, 10(4), p. 181. doi: 10.4103/JPBS.JPBS\_126\_18.
- Chandradevi, W. A. et al. (2018) ‘Pemanfaatan Daun Kelor (Moringa oleifera) Terhadap Pemulihan Struktur Pankreas Mencit Diabetik’, *Biotropic : The Journal of Tropical Biology*, 2(2). doi: 10.29080/biotropic.2018.2.2.85-92.
- Cho, N. H. et al. (2018) ‘IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045’, *Diabetes Research and Clinical Practice*, 138. doi: 10.1016/j.diabres.2018.02.023.
- Citra Rani, K. and Deasy Rosita Dewi, A. (2019) ‘MODUL PELATIHAN KANDUNGAN NUTRISI TANAMAN KELOR Disusun oleh’.
- Daba, M.-H., Elmasry, A. and El-Karef, A. A. (2015) *Effect of moringa oleifera with and without metformin on an experimental model of metabolic syndrome in rats*, *International Journal of Advanced Research*. Available at:

- <http://www.journalijar.com> (Accessed: 26 May 2021).
- David G. Gardner, MD, M. and Dolores Shoback, M. (2011) *Greenspan's Basic & Clinical Endocrinology, Ninth Edition, Greenspan's Basic & Clinical Endocrinology*.
- Diabetes Risk Factors / CDC* (2021). Available at:  
<https://www.cdc.gov/diabetes/basics/risk-factors.html> (Accessed: 10 August 2021).
- El-Desouki, N. I. and Basyony, M. A. (2015) *Moringa oleifera leaf extract ameliorates glucose, insulin and pancreatic beta cells disorder in alloxan-induced diabetic rats.*, *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. Available at:  
[https://www.rjpbcos.com/pdf/2015\\_6\(3\)/\[90\].pdf](https://www.rjpbcos.com/pdf/2015_6(3)/[90].pdf) (Accessed: 12 September 2021).
- Eroschenko, V. P. et al. (2012) ‘Atlas Histologi ffifffWffifwww’, *Atlas Histologi diFiore dengan Korelasi Fungsional*, 11.
- Furman, B. L. (2021) ‘Streptozotocin-Induced Diabetic Models in Mice and Rats’, *Current Protocols*, 1(4), p. e78. doi: 10.1002/CPZ1.78.
- Galicia-Garcia, U. et al. (2020) ‘Pathophysiology of type 2 diabetes melitus’, *International Journal of Molecular Sciences*. doi: 10.3390/ijms21176275.
- Gopalakrishnan, L., Doriya, K. and Kumar, D. S. (2016) ‘Moringa oleifera: A review on nutritive importance and its medicinal application’, *Food Science and Human Wellness*, 5(2), pp. 49–56. doi: 10.1016/J.FSHW.2016.04.001.
- Gupta, R. et al. (2011) ‘Antidiabetic and antioxidant potential of β-sitosterol in streptozotocin-induced experimental hyperglycemia’, *Journal of Diabetes*, 3(1). doi: 10.1111/j.1753-0407.2010.00107.x.
- Hajiaghaalipour, F. et al. (2015) ‘Modulation of glucose transporter protein by dietary flavonoids in type 2 diabetes melitus’, *International Journal of Biological Sciences*, 11(5). doi: 10.7150/ijbs.11241.
- Halban, P. A. et al. (2014) ‘β-Cell failure in type 2 diabetes: Postulated mechanisms and prospects for prevention and treatment’, *Journal of Clinical Endocrinology and Metabolism*, 99(6). doi: 10.1210/jc.2014-1425.
- Hariyati, R. T. S. (2010) ‘Mengenal Systematic Review Theory dan Studi Kasus’, *Jurnal Keperawatan Indonesia*, 13(2). doi: 10.7454/jki.v13i2.242.
- Ighodaro, O. M., Adeosun, A. M. and Akinloye, O. A. (2017) ‘Alloxan-induced diabetes, a common model for evaluating the glycemic-control potential of therapeutic compounds and plants extracts in experimental studies’, *Medicina (Lithuania)*. doi: 10.1016/j.medici.2018.02.001.

- International Diabetes Federation (2020) *Members*. Available at:  
<https://idf.org/our-network/regions-members/western-pacific/members/104-indonesia.html> (Accessed: 20 April 2021).
- ITIS Standard Report Page: Moringa oleifera* (2021). Available at:  
[https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=503874#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=503874#null) (Accessed: 10 July 2021).
- Jomová, K. et al. (2019) ‘A switch between antioxidant and prooxidant properties of the phenolic compounds myricetin, morin, 3’,4’-dihydroxyflavone, taxifolin and 4-hydroxy-coumarin in the presence of copper(II) ions: A spectroscopic, absorption titration and DNA damage study’, *Molecules*, 24(23). doi: 10.3390/molecules24234335.
- Kasolo, J. N. et al. (2010) ‘Phytochemicals and uses of Moringa oleifera leaves in Ugandan rural communities’, *Journal of Medicinal Plants Research*, 4(9). doi: 10.5897/JMPR10.492.
- Kemenkes RI (2014) ‘Info Datin Pusat Data dan Informasi Kementerian Kesehatan RI, Situasi dan Analisis DIABETES’, *Pusat Data dan Informasi Kementerian RI*.
- Kerimi, A. and Williamson, G. (2016) ‘At the interface of antioxidant signalling and cellular function: Key polyphenol effects’, *Molecular nutrition & food research*. doi: 10.1002/mnfr.201500940.
- Kerner, A. W. and Brückel, J. (2014) ‘Definition, Classification and Diagnosis of Diabetes Melitus’, *Exp Clin Endocrinol Diabetes*, 122, pp. 384–386. doi: 10.1055/s-0034-1366278.
- Khalid Al-Ishaq, R. et al. (2019) ‘biomolecules Flavonoids and Their Anti-Diabetic Effects: Cellular Mechanisms and Effects to Improve Blood Sugar Levels’. doi: 10.3390/biom9090430.
- King, A. J. (2012) ‘The use of animal models in diabetes research’, *British Journal of Pharmacology*, 166(3), p. 877. doi: 10.1111/J.1476-5381.2012.01911.X.
- Kumar, V., Abbas, A. K. and Aster, J. C. (2018) *Robbins Basic Pathology, Tenth Edition*, Elsevier Inc.
- Larantukan, S. V. M. et al. (2014) ‘Pemberian Ekstrak Etanol Kulit Batang Kelor Glukosa Darah Tikus Hiperglikemia’, *Indonesia Medicus Veterinus*, 3(4).
- Liang, L. et al. (2019) ‘Nutritional compositions of Indian Moringa oleifera seed and antioxidant activity of its polypeptides’, *Food Science and Nutrition*, 7(5). doi: 10.1002/fsn3.1015.
- López, M. et al. (2016) ‘Hypothalamic AMPK: A canonical regulator of whole-

- body energy balance', *Nature Reviews Endocrinology*. doi: 10.1038/nrendo.2016.67.
- Luippold, G. et al. (2016) 'Short- and Longterm Glycemic Control of Streptozotocin-Induced Diabetic Rats Using Different Insulin Preparations', *PLOS ONE*, 11(6), p. e0156346. doi: 10.1371/JOURNAL.PONE.0156346.
- Mahmoud, A. M. et al. (2019) 'Beneficial effects of citrus flavonoids on cardiovascular and metabolic health', *Oxidative Medicine and Cellular Longevity*. doi: 10.1155/2019/5484138.
- Moher, D. et al. (2016) 'Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement', *Revista Espanola de Nutricion Humana y Dietetica*, 20(2). doi: 10.1186/2046-4053-4-1.
- Mussbacher, M. et al. (2019) 'Cell type specific roles of nf-kb linking inflammation and thrombosis', *Frontiers in Immunology*, 10(FEB). doi: 10.3389/FIMMU.2019.00085/FULL.
- Nada, S. et al. (2015) 'Evaluation Of Moringa Oleifera Leaves Extract Effects On Streptozotocin-Induced Diabetic Rats', *Advances in Food Sciences*, 37, p. 10. Available at: [https://scholar.cu.edu.eg/sites/default/files/abbasms/files/evaluation\\_of\\_moringa\\_oleifera\\_leaves\\_extract\\_effects\\_on\\_streptozotocin-induced\\_diabetic\\_rats.pdf](https://scholar.cu.edu.eg/sites/default/files/abbasms/files/evaluation_of_moringa_oleifera_leaves_extract_effects_on_streptozotocin-induced_diabetic_rats.pdf) (Accessed: 17 September 2021).
- Nagaki, Y., Ito, K. and Kuwahara, M. (2016) 'WTC rat has unique characteristics such as resistant to streptozotocin', *Biochemistry and Biophysics Reports*, 8, pp. 157–161. doi: 10.1016/J.BBREP.2016.08.024.
- Naik, R. S. et al. (2019) 'Studies on the effect of moringa leaf extract on blood glucose and insulin levels in streptozotocin induced diabetic rats', ~ 189 ~ *The Pharma Innovation Journal*, 8(11), pp. 189–194. Available at: <http://www.thepharmajournal.com> (Accessed: 17 September 2021).
- Nova, E. et al. (2020) 'Potential of moringa oleifera to improve glucose control for the prevention of diabetes and related metabolic alterations: A systematic review of animal and human studies', *Nutrients*. doi: 10.3390/nu12072050.
- Omeodu, S. I., Onyegeme-Okerenta and Nnake, B. M. (2017) 'Evaluation Of Anti-Diabetic And Anti-Lipidaemic Potential Of Aqueous Leaf Extract Of Moringa Oleifera On Streptozotocin Diabetic-Induced Albino Rats', *European Journal of Advanced Research in Biological and Life Sciences*, 5(2). Available at: [www.idpublications.org](http://www.idpublications.org).
- Omodanisi, E. I. et al. (2017) 'Assessment of the anti-hyperglycaemic, anti-inflammatory and antioxidant activities of the methanol extract of moringa oleifera in diabetes-induced nephrotoxic male wistar rats', *Molecules*, 22(4). doi: 10.3390/molecules22040439.

- Oyeyinka, A. T. and Oyeyinka, S. A. (2018) 'Moringa oleifera as a food fortificant: Recent trends and prospects', *Journal of the Saudi Society of Agricultural Sciences*, 17(2), pp. 127–136. doi: 10.1016/J.JSSAS.2016.02.002.
- Panche, A. N., Diwan, A. D. and Chandra, S. R. (2016) 'Flavonoids: an overview', *Journal of Nutritional Science*, 5, pp. 1–15. doi: 10.1017/jns.2016.41.
- Rachmawani, N. R. and Oktarlina, R. Z. (2017) *Khasiat Pemberian Buncis (Phaseolus vulgaris L.) sebagai Terapi Alternatif Diabetes Melitus Tipe 2, Jurnal Majority*.
- Renée A. Street, Gerhard Prinsloo, L. J. M. (2017) 'Alkaloids potential health Benefits and toxicity', *Utilisation of Bioactive Compounds from Agricultural and Food Waste*, pp. 60–85. doi: 10.1201/9781315151540-3.
- Riskesdas (2013) *PEMANFAATAN PENGOBATAN TRADISIONAL*. Available at: <https://dinkes.sarolangunkab.go.id/berita-pemanfaatan-pengobatan-tradisional.html> (Accessed: 18 May 2021).
- Rizkayanti, R., Diah, A. W. M. and Jura, M. R. (2017) 'Uji Aktivitas Antioksidan Ekstrak Air dan Ekstrak Etanol Daun Kelor (Moringa Oleifera LAM)', *Jurnal Akademika Kimia*, 6(2). doi: 10.22487/j24775185.2017.v6.i2.9244.
- Rockwood, J., Anderson, B. and Casamatta, D. (2013) 'Potential Uses Of Moringa Oleifera And An Examination Of Antibiotic Efficacy Conferred By M. Oleifera Seed And Leaf Extracts Using Crude Extraction Techniques Available To Underserved Indigenous Populations', *undefined*.
- Satyarsa, A. B. S. (2019) 'Potential Effects of Alkaloid vindolicine Substances in Tapak Dara Leafs (Catharanthus roseus (L.) G. Don) in Reducing Blood Glucose Levels', *Journal of Medicine and Health*, 2(4). doi: 10.28932/jmh.v2i4.1057.
- Sekar, V. et al. (2019) 'Mangiferin from Mangifera indica fruits reduces post-prandial glucose level by inhibiting  $\alpha$ -glucosidase and  $\alpha$ -amylase activity', *South African Journal of Botany*, 120. doi: 10.1016/j.sajb.2018.02.001.
- Semaan, D. G. et al. (2017) 'In vitro anti-diabetic activity of flavonoids and pheophytins from Allophylus cominia Sw. on PTP1B, DPPIV, alpha-glucosidase and alpha-amylase enzymes', *Journal of Ethnopharmacology*, 203. doi: 10.1016/j.jep.2017.03.023.
- Shamseer, L. et al. (2015) 'Preferred reporting items for systematic review and meta-analysis protocols (prisma-p) 2015: Elaboration and explanation', *BMJ (Online)*. doi: 10.1136/bmj.g7647.
- Sherwood, L. (2014) 'Human Anatomy and Physiology from Cell to System', in *Fisiologi Manusia dari Sel ke Sistem*.

- Shewasinad, A. *et al.* (2019) ‘Antidiabetic Activity of Methanol Extract and Fractions of Thymus schimperi Ronniger Leaves in Normal and Streptozotocin Induce Diabetic Mice’, *Iranian Journal of ...*, 2019(October).
- Shih, M. C. *et al.* (2011) ‘Effect of different parts (leaf, stem and stalk) and seasons (summer and winter) on the chemical compositions and antioxidant activity of Moringa oleifera’, *International Journal of Molecular Sciences*, 12(9). doi: 10.3390/ijms12096077.
- Siswanto, S. (2012) ‘SYSTEMATIC REVIEW SEBAGAI METODE PENELITIAN UNTUK MENSINTESIS HASIL-HASIL PENELITIAN (SEBUAH PENGANTAR)’, *Buletin Penelitian Sistem Kesehatan*, 13(4). doi: 10.22435/bpsk.v13i4.
- Smirnov, A. V., Snigur, G. L. and Voronkova, M. P. (2012) ‘Pancreatic Islet Beta-Cell Apoptosis in Experimental Diabetes Melitus’, *Apoptosis and Medicine*. doi: 10.5772/51411.
- Soebagijo Adi Soelistijo, Novida, H. and Rudijanto, A. (2015) *PERKENI 2015, Perkeni*.
- Soelistijo, S. A. *et al.* (2015) *Konsensus Pengendalian dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2015, Perkeni*. doi: 10.1017/CBO9781107415324.004.
- Speight, J. and Pouwer, F. (2021) ‘Diabetes Melitus Type 1’, *Cambridge Handbook of Psychology, Health and Medicine: Third Edition*, pp. 477–480. doi: 10.29309/tpmj/2017.24.12.614.
- Sudha, R., Chandra Philip, X. and Suriyakumari, K. (2020) ‘Phytochemical Constituents of Leaves of Moringa oleifera Grow in Cuddalore District, Tamil Nadu, India’, *Clinical and Applied Health Science*. doi: 10.5005/jp-journals-10082-02270.
- Sulastri, E. *et al.* (2018) ‘Total phenolic, total flavonoid, quercetin content and antioxidant activity of standardized extract of moringa oleifera leaf from regions with different elevation’, *Pharmacognosy Journal*, 10(6). doi: 10.5530/pj.2018.6s.20.
- Sulistyorini, R. *et al.* (2015) ‘Pengaruh Ekstrak Etanol Daun Kelor (Moringa oleifera) pada Ekspresi Insulin dan Insulitis Tikus Diabetes Melitus’, *Majalah Kedokteran Bandung*, 47(2). doi: 10.15395/MKB.V47N2.456.
- Tahir, M., Hikmah, N. and Rahmawati, R. (2016) ‘Analisis Kandungan Vitamin C Dan B- Karoten Dalam Daun Kelor (Moringa Oleifera Lam.) Dengan Metode Spektrofotometri Uv–Vis’, *Jurnal Fitofarmaka Indonesia*, 3(1). doi: 10.33096/jffi.v3i1.173.
- Tilong, A. D. (2012) ‘Kelor Penakluk Diabetes’, *Penerbit Diva press. Yogyakarta*.

- Tochukwu, N., Monago, C. and Chuku, L. (2016) 'Histological Effect of Combined Ethanol Extract of Moringa oleifera and Pleurotus ostreatus on the Pancreas of Alloxan-induced Diabetic Wistar Albino Rats', *Journal of Applied Life Sciences International*, 5(3). doi: 10.9734/jalsi/2016/26108.
- Tundis, R. et al. (2016) 'Poncirus trifoliata (L.) Raf.: Chemical composition, antioxidant properties and hypoglycaemic activity via the inhibition of  $\alpha$ -amylase and  $\alpha$ -glucosidase enzymes', *Journal of Functional Foods*, 25. doi: 10.1016/j.jff.2016.06.034.
- Wei, Q. et al. (2020) 'Pathological Mechanisms in Diabetes of the Exocrine Pancreas: What's Known and What's to Know', *Frontiers in Physiology*, 11. doi: 10.3389/FPHYS.2020.570276.
- WHO (2021) *Diabetes*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes> (Accessed: 20 April 2021).
- Yan, Y. et al. (2020) 'Use of Chlorogenic Acid against Diabetes Melitus and Its Complications', *Journal of Immunology Research*. doi: 10.1155/2020/9680508.
- Yassa, H. D. and Tohamy, A. F. (2014) 'Extract of Moringa oleifera leaves ameliorates streptozotocin-induced Diabetes melitus in adult rats', *Acta Histochemica*, 116(5). doi: 10.1016/j.acthis.2014.02.002.
- Zhong, F. and Jiang, Y. (2019) 'Endogenous Pancreatic  $\beta$  Cell Regeneration: A Potential Strategy for the Recovery of  $\beta$  Cell Deficiency in Diabetes', *Frontiers in Endocrinology*, 10(FEB), p. 101. doi: 10.3389/FENDO.2019.00101.
- Zhou, Q. and Melton, D. A. (2018) 'Pancreas regeneration', *Nature*, 557(7705), p. 351. doi: 10.1038/S41586-018-0088-0.