

DAFTAR PUSTAKA

- Alharbi, M. A. (2020) ‘Is low serum albumin a predictor sign of the incidence of peritoneal dialysis-associated peritonitis? a quasi-systematic review’, *Saudi Journal of Kidney Diseases and Transplantation*, 31(2), pp. 320–334. doi: 10.4103/1319-2442.284006.
- Andreoli, M. C. C., dan Totoli, C. (2020) ‘Peritoneal dialysis’, *Revista da Associacao Medica Brasileira*, 66, pp. 37–44. doi: 10.1590/1806-9282.66.S1.37.
- Batubara, S. O. (2012) *Analisis Faktor Resiko terhadap Komplikasi CAPD di RSUP Dr. Moewardi Surakarta dan RSUP Dr. Sardjito Yogyakarta*, Tesis Program Pasca Sarjana. Universitas Indonesia.
- Chang, H. H. dkk. (2021) ‘Diabetes Is the Most Critical Risk Factor of Adverse Complications After Peritoneal Dialysis Catheter Placement’, *Frontiers in Medicine*, 8(October), pp. 1–9. doi: 10.3389/fmed.2021.719345.
- Cheikh Hassan, H. I. dkk. (2022) ‘Association of Peritonitis With Cardiovascular Mortality Over Time in the Peritoneal Dialysis Population: An Australia and New Zealand Dialysis and Transplant Registry Study’, *Kidney International Reports*, 7(11), pp. 2388–2396. doi: 10.1016/j.ekir.2022.08.008.
- Chen, H. L. dkk. (2019) ‘Risk factors associated with outcomes of peritoneal dialysis in Taiwan: An analysis using a competing risk model’, *Medicine (United States)*, 98(6). doi: 10.1097/MD.00000000000014385.
- Dao Bui Quy, Q. dkk. (2020) ‘Overhydration and low serum prealbumin predict peritoneal dialysis-related peritonitis in continuous ambulatory peritoneal dialysis patients’, *BMC Nephrology*, 21(1), pp. 1–8. doi: 10.1186/s12882-020-02178-w.
- Davies, S. J. dkk. (2021) ‘Low Serum Potassium Levels and Clinical Outcomes in Peritoneal Dialysis—International Results from PDOPPS’, *Kidney International Reports*, 6(2), pp. 313–324. doi: 10.1016/j.ekir.2020.11.021.
- Dehghani, A. dkk. (2022) ‘Prevalence of chronic kidney diseases and its determinants among Iranian adults: results of the first phase of Shahedieh cohort study’, *BMC Nephrology*, 23(1), pp. 1–11. doi: 10.1186/s12882-022-02832-5.
- van Eck van der Sluijs, A. dkk. (2021) ‘Bleeding risk of haemodialysis and peritoneal dialysis patients’, *Nephrology Dialysis Transplantation*, 36(1), pp. 170–175. doi: 10.1093/NDT/GFAA216.

- He, P. dkk. (2020) ‘Red blood cell distribution width and peritoneal dialysis-associated peritonitis prognosis’, *Renal Failure*, 42(1), pp. 613–621. doi: 10.1080/0886022X.2020.1786401.
- Htay, H. dkk. (2019) ‘Comparison of clinical presentation and outcomes of peritonitis in the elderly and younger peritoneal dialysis patients’, *Peritoneal Dialysis International*, 39(2), pp. 163–168. doi: 10.3747/pdi.2018.00056.
- Hu, J., Zhang, H., dan Yi, B. (2021) ‘Peritoneal transport status and first episode of peritonitis: a large cohort study’, *Renal Failure*, 43(1), pp. 1094–1103. doi: 10.1080/0886022X.2021.1949350.
- Huo, Z. dkk. (2022) ‘Hypokalemia Duration in the First Year Associated with Subsequent Peritoneal Dialysis-Associated Peritonitis: A Multicenter Retrospective Cohort Study’, *Journal of Clinical Medicine*, 11(24). doi: 10.3390/jcm11247518.
- Hwang, S. D. dkk. (2019) ‘Impact of body mass index on survival in patients undergoing peritoneal dialysis: Analysis of data from the Insan Memorial End-Stage Renal Disease Registry of Korea (1985–2014)’, *Kidney Research and Clinical Practice*, 38(2), pp. 239–249. doi: 10.23876/j.krcp.18.0106.
- Indonesian Renal Registry (2018) *11th report Of Indonesian renal registry 2018, Indonesian Renal Registry (IRR)*.
- Indrayanti, S. dkk. (2019) ‘Risk Factors for Chronic Kidney Disease: A Case-Control Study in a District Hospital in Indonesia’, *Journal of Pharmaceutical Sciences and Research*, 11(7), pp. 2549–2554. Available at: <http://sipeg.univpancasila.ac.id/uploads/repository/lampiran/DokumenLampiran-18082020093419.pdf>.
- Jameson, L. J. dan Loscalzo, J. (2017) *Harrison’s Nephrology and Acid-Base Disorder*. 3rd edn. New York: McGraw-Hill Education.
- Janice L., H. dan Cheever, K. H. (2018) *Brunner & Suddarth’s Textbook of Medical - Surgical Nursing*. 14th edn. Philadelphia: Wolters Kluwer.
- Jegatheesan, D. dkk. (2018) ‘The relationship between body mass index and organism-specific peritonitis’, *Peritoneal Dialysis International*, 38(3), pp. 206–214. doi: 10.3747/pdi.2017.00188.
- Jones, T. (2015) *Renal and Urinary Systems*. 4th edn. New York: Elsevier.
- Kang, M. dkk. (2021) ‘Evolving outcomes of peritoneal dialysis: Secular trends at a single large center over three decades’, *Kidney Research and Clinical Practice*, 40(3), pp. 472–483. doi: 10.23876/j.krcp.21.020.

Kemenkes RI (2018) *Laporan Riskesdas 2018 Kementerian Kesehatan Republik Indonesia, Laporan Nasional Riskesdas 2018*. Jakarta. Available at: <http://www.yankeks.kemkes.go.id/assets/downloads/PMK No. 57 Tahun 2013 tentang PTRM.pdf>.

Kerschbaum, J., König, P., dan Rudnicki, M. (2012) ‘Risk factors associated with peritoneal-dialysis-related peritonitis’, *International Journal of Nephrology*, 2012(April). doi: 10.1155/2012/483250.

Kojima, S. dkk. (2021) ‘Evaluation of modifiable risk factors for peritoneal dialysis-related peritonitis: a single-center retrospective cohort study’, *Renal Replacement Therapy*, 7(1), pp. 1–7. doi: 10.1186/s41100-021-00343-x.

Kovesdy, C. P. (2022) ‘Epidemiology of chronic kidney disease: an update 2022’, *Kidney International Supplements*, 12(1), pp. 7–11. doi: 10.1016/j.kisu.2021.11.003.

Li, P. K. T. dkk. (2022) ‘ISPD peritonitis guideline recommendations: 2022 update on prevention and treatment’, *Peritoneal Dialysis International*, 42(2), pp. 110–153. doi: 10.1177/08968608221080586.

Lin, Z. dkk. (2020) ‘Establishment of age group classification for risk stratification in glioma patients’, *BMC Neurology*, 20(1), pp. 1–11. doi: 10.1186/s12883-020-01888-w.

Liu, D. dkk. (2021) ‘Degree and duration of hypokalemia associated with peritonitis in patients undergoing peritoneal dialysis’, *International Journal of Clinical Practice*, 75(8), pp. 1–12. doi: 10.1111/ijcp.14188.

Ma, X. dkk. (2020) ‘Analysis of risk factors and outcome in peritoneal dialysis patients with early-onset peritonitis: A multicentre, retrospective cohort study’, *BMJ Open*, 10(2), pp. 1–9. doi: 10.1136/bmjopen-2019-029949.

Maemun, S. dkk. (2020) ‘Is hypoalbuminemia a predictor marker of mortality?’, *Health Science Journal of Indonesia*, 11(2), pp. 121–125. doi: 10.22435/hsji.v11i2.3072.

Masturoh, I. dan Anggita T., N. (2018) *Metodologi Penelitian Kesehatan*. Jakarta: Kemenkes RI.

McAninch, J. W. dan Lue, T. F. (2020) *Smith & Tanagho’s General Urology*. Nineteenth. San Fransisco: McGraw-Hill Education.

Molitoris, B. A. dkk. (2022) ‘Albumin Uptake and Processing By the Proximal Tubule: Physiological, Pathological, and Therapeutic Implications’, *Physiological Reviews*, 102(4), pp. 1625–1667. doi: 10.1152/physrev.00014.2021.

- Musoke, J. dkk. (2020) ‘The prevalence and bacterial distribution of peritonitis amongst adults undergoing continuous ambulatory peritoneal dialysis at Universitas hospital’, *Southern African Journal of Infectious Diseases*, 35(1), pp. 1–5. doi: 10.4102/sajid.v35i1.104.
- Nadeau-Fredette, A. C. dkk. (2016) ‘Center-specific factors associated with peritonitis risk—a multi-center registry analysis’, *Peritoneal Dialysis International*, 36(5), pp. 509–518. doi: 10.3747/pdi.2015.00146.
- Obi, Y. dkk. (2018) ‘Impact of Obesity on Modality Longevity, Residual Kidney Function, Peritonitis, and Survival Among Incident Peritoneal Dialysis Patients’, *Am J Kidneys Dis*, 71(6), pp. 802–813. doi: 10.1053/j.gastro.2016.08.014.CagY.
- Perl, J. dkk. (2020) ‘Peritoneal Dialysis–Related Infection Rates and Outcomes: Results From the Peritoneal Dialysis Outcomes and Practice Patterns Study (PDOPPS)’, *American Journal of Kidney Diseases*, 76(1), pp. 42–53. doi: 10.1053/j.ajkd.2019.09.016.
- Piraino, B. M. (2022) ‘Correction of Hypokalemia in Peritoneal Dialysis Patients May Decrease Peritonitis Risk’, *American Journal of Kidney Diseases*, 80(5), pp. 564–565. doi: 10.1053/j.ajkd.2022.06.001.
- Prasad, R., Jha, R. K. dan Keerti, A. (2022) ‘Chronic Kidney Disease: Its Relationship With Obesity’, *Cureus*, 14(10). doi: 10.7759/cureus.30535.
- Purwati, S. (2018) ‘Analisis Faktor Risiko Penyebab Kejadian Penyakit Gagal Ginjal Kronik (GGK) Di Ruang Hemodialisa RS Dr. Moewardi’, (*Jkg*) *Jurnal Keperawatan Global*, 3(1), pp. 15–27. doi: 10.37341/jkg.v3i1.44.
- Quero, M. dkk. (2021) ‘Impact of obesity on the evolution of outcomes in peritoneal dialysis patients’, *Clinical Kidney Journal*, 14(3), pp. 969–982. doi: 10.1093/ckj/sfaa055.
- Shi, Y. dkk. (2022) ‘Low Serum Albumin Is Associated with Poor Prognosis in Patients Receiving Peritoneal Dialysis Treatment’, *Journal of Healthcare Engineering*, 2022. doi: 10.1155/2022/7660806.
- Soesilo, T. D. (2019) *Ragam dan Prosedur Penelitian Tindakan*.
- Song, P. dkk. (2022) ‘Microbiology and Outcome of Peritoneal Dialysis-Related Peritonitis in Elderly Patients: A Retrospective Study in China’, *Frontiers in Medicine*, 9(April), pp. 1–9. doi: 10.3389/fmed.2022.799110.
- Sukul, N. dkk. (2019) ‘Patient-reported advantages and disadvantages of peritoneal dialysis: Results from the PDOPPS’, *BMC Nephrology*, 20(1), pp. 1–10. doi: 10.1186/s12882-019-1304-3.

- Supono (2008) *Faktor - Faktor yang Berkontribusi Terjadinya Peritonitis pada Pasien Continuous Ambulatory Peritoneal Dialysis di Rumah Sakit Umum Dr. Saiful Anwar Malang*. Universitas Indonesia.
- Tao, W. dkk. (2022) ‘Peritoneal-scrotal dialysate leakage demonstrated on SPECT/CT imaging in a patient on peritoneal dialysis’, *Radiology Case Reports*, 17(8), pp. 2643–2646. doi: 10.1016/j.radcr.2022.05.003.
- Tekkarişmaz, N. dan Torun, D. (2020) ‘Long-term clinical outcomes of peritoneal dialysis patients: 9-year experience of a single centre in Turkey’, *Turkish Journal of Medical Sciences*, 50(2), pp. 386–397. doi: 10.3906/sag-1909-98.
- Tian, Y. dkk. (2017) ‘Risk factors and outcomes of early-onset peritonitis in Chinese peritoneal dialysis patients’, *Kidney and Blood Pressure Research*, 42(6), pp. 1266–1276. doi: 10.1159/000485930.
- Ueda, R. dkk. (2019) ‘Effect of diabetes on incidence of peritoneal dialysis-associated peritonitis’, *PLoS ONE*, 14(12), pp. 1–10. doi: 10.1371/journal.pone.0225316.
- Ueki, K. dkk. (2022) ‘A case of peritoneal dialysis-related peritonitis caused by dialysate leakage with successful treatment by intravenous and intraperitoneal antibiotic therapy’, *CEN case reports*, 11(2), pp. 161–165. doi: 10.1007/s13730-021-00644-4.
- Urden, L. D., Stacy, K. M., dan Lough, M. E. (2021) *Critical Care Nursing: Diagnosis and Management*. Ninth. Missouri: Elsevier. doi: 10.4037/ccn2017185.
- Vaidya, S. R. dan Aedula, N. R. (2022) *Chronic Renal Failure, StatPearls Publishing*.
- Vaios, V. dkk. (2019) ‘Assessment and management of hypertension among patients on peritoneal dialysis’, *Clinical Journal of the American Society of Nephrology*, 14(2), pp. 297–305. doi: 10.2215/CJN.07480618.
- Wang, H. dkk. (2018) ‘Risk factors for peritoneal dialysis-associated peritonitis’, *European Journal of Inflammation*, 16. doi: 10.1177/2058739218772243.
- Wilson, S. dkk. (2021) ‘Chronic kidney disease: Definition, updated epidemiology, staging, and mechanisms of increased cardiovascular risk’, *Journal of Clinical Hypertension*, 23(4), pp. 831–834. doi: 10.1111/jch.14186.
- Wu, H. dkk. (2020) ‘Incidence and risk factors of peritoneal dialysis-related peritonitis in elderly patients: A retrospective clinical study’, *Peritoneal Dialysis International*, 40(1), pp. 26–33. doi: 10.1177/0896860819879868.

- Yamada, S. dan Inaba, M. (2021) ‘Potassium metabolism and management in patients with ckd’, *Nutrients*, 13(6), pp. 1–19. doi: 10.3390/nu13061751.
- Yanti, S. (2021) ‘Faktor Risiko Peritonitis Pada Pasien Continuous Ambulatory Peritoneal Dialysis (Capd) Literature Review’, *Keperawatan FIK Unsyiah*. Available at: http://digilib.unisyah.ac.id/5947/1/SRI_YANTI_1910201220_KEPERAWATAN_S1_Naspub - sri yanti.pdf.
- Zeng, Y. dkk. (2020) ‘The influence of seasonal factors on the incidence of peritoneal dialysis-associated peritonitis’, *Renal Failure*, 42(1), pp. 807–817. doi: 10.1080/0886022X.2020.1804401.