

DAFTAR PUSTAKA

- Abdissa, D. and Hirpa, D. (2022) ‘Dyslipidemia and its associated factors among adult diabetes outpatients in West Shewa zone public hospitals, Ethiopia’, *BMC Cardiovascular Disorders*, 22(1), p. 39. Available at: <https://doi.org/10.1186/s12872-022-02489-w>.
- Abdul Basit, K. *et al.* (2023) ‘Changes in the prevalence of diabetes, prediabetes and associated risk factors in rural Baluchistan; a secondary analysis from repeated surveys (2002–2017)’, *PLOS ONE*, 18(4), p. e0284441. Available at: <https://doi.org/10.1371/journal.pone.0284441>.
- ADA (2024) *Alcohol and Diabetes, Health-Wellness*.
- Aldossari, K.K. *et al.* (2018) ‘Prevalence of Prediabetes, Diabetes, and Its Associated Risk Factors among Males in Saudi Arabia: A Population-Based Survey.’, *Journal of diabetes research*, 2018, p. 2194604. Available at: <https://doi.org/10.1155/2018/2194604>.
- Alkandari, A. *et al.* (2020) ‘Adult Diabetes and Prediabetes Prevalence in Kuwait: Data from the Cross-Sectional Kuwait Diabetes Epidemiology Program’, *Journal of Clinical Medicine*, 9(11), p. 3420. Available at: <https://doi.org/10.3390/jcm9113420>.
- Al-Mohaissen, M.A., Lee, T. and Alamri, A.F. (2020) ‘Vitamin D levels, prediabetes risk and hemoglobin A1c levels in young non-diabetic Saudi women.’, *Journal of diabetes investigation*, 11(5), pp. 1344–1351. Available at: <https://doi.org/10.1111/jdi.13226>.
- Alomari, A. and Al Hisnah, S. (2022) ‘Prevalence of Prediabetes and Associated Risk Factor Assessment Among Adults Attending Primary Healthcare Centers in Al Bahah, Saudi Arabia: A Cross-Sectional Study’, *Cureus* [Preprint]. Available at: <https://doi.org/10.7759/cureus.29465>.
- Alvarez, S. *et al.* (2024) *Prediabetes*.
- American Diabetes Association (2024) *Genetics of Diabetes*. Available at: <https://diabetes.org/about-diabetes/genetics-diabetes> (Accessed: 6 October 2024).
- American Heart Association (2024) *Diabetes Risk Factors*. Available at: <https://www.heart.org/en/health-topics/diabetes/understand-your-risk-for-diabetes> (Accessed: 6 October 2024).
- Ampeire, I.P., Kawugezi, P.C. and Mulogo, E.M. (2023) ‘Prevalence of prediabetes and associated factors among community members in rural Isingiro district.’,

- BMC public health*, 23(1), p. 958. Available at: <https://doi.org/10.1186/s12889-023-15802-9>.
- Al Amri, T. et al. (2019) ‘The Association Between Prediabetes and Dyslipidemia Among Attendants of Primary Care Health Centers in Jeddah, Saudi Arabia’, *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, Volume 12, pp. 2735–2743. Available at: <https://doi.org/10.2147/DMSO.S233717>.
- Andes, L.J. et al. (2020) ‘Prevalence of Prediabetes Among Adolescents and Young Adults in the United States, 2005–2016.’, *JAMA pediatrics*, 174(2), p. e194498. Available at: <https://doi.org/10.1001/jamapediatrics.2019.4498>.
- Andriani, R.N. et al. (2022) ‘Analysis of Prediabetes Risk Factors at Primary Health Care Centers’, *Review of Primary Care Practice and Education (Kajian Praktik dan Pendidikan Layanan Primer)*, 5(2), p. 52. Available at: <https://doi.org/10.22146/rpcpe.74552>.
- Apidechkul, T. et al. (2022) ‘Epidemiology of prediabetes mellitus among hill tribe adults in Thailand’, *PLOS ONE*, 17(7), p. e0271900. Available at: <https://doi.org/10.1371/journal.pone.0271900>.
- Artese, A., Stamford, B.A. and Moffatt, R.J. (2019) ‘Cigarette Smoking: An Accessory to the Development of Insulin Resistance.’, *American journal of lifestyle medicine*, 13(6), pp. 602–605. Available at: <https://doi.org/10.1177/1559827617726516>.
- Asriati, A. (2023) ‘Analisis Perilaku Konsumsi Makanan Dan Minuman Manis Terhadap Prediabetes Remaja Di Kota Jayapura’, *Preventif: Jurnal Kesehatan Masyarakat*, 14(3), pp. 495–511. Available at: <https://doi.org/10.22487/preventif.v14i3.970>.
- Astuti, A. (2019) ‘Usia, Obesitas dan Aktifitas Fisik Beresiko Terhadap Prediabetes’, *Jurnal Endurance*, 4(2), p. 319. Available at: <https://doi.org/10.22216/jen.v4i2.3757>.
- Bahadoran, Z. et al. (2023) ‘Higher daily physical activity levels may facilitate pre-diabetes regression to normoglycemia: A longitudinal study among an Iranian population.’, *Preventive medicine reports*, 34, p. 102233. Available at: <https://doi.org/10.1016/j.pmedr.2023.102233>.
- Bell, K. et al. (2020) ‘A position statement on screening and management of prediabetes in adults in primary care in Australia’, *Diabetes Research and Clinical Practice*, 164, p. 108188. Available at: <https://doi.org/10.1016/j.diabres.2020.108188>.
- Berberich, A.J. and Hegele, R.A. (2022) ‘A Modern Approach to Dyslipidemia’, *Endocrine Reviews*, 43(4), pp. 611–653. Available at: <https://doi.org/10.1210/endrev/bnab037>.

- BKPK Kemenkes (2024) *Survei Kesehatan Indonesia (SKI) 2023 dalam Angka*.
- Block, G. et al. (2015) ‘Diabetes Prevention and Weight Loss with a Fully Automated Behavioral Intervention by Email, Web, and Mobile Phone: A Randomized Controlled Trial Among Persons with Prediabetes’, *Journal of Medical Internet Research*, 17(10), p. e240. Available at: <https://doi.org/10.2196/jmir.4897>.
- BPS (2024) *Jumlah Penduduk Pertengahan Tahun (Ribu Jiwa), 2022-2024*. Available at: <https://www.bps.go.id/id/statistics-table/2/MTk3NSMy/jumlah-penduduk-pertengahan-tahun--ribu-jiwa-.html> (Accessed: 5 January 2025).
- Brannick, B. and Dagogo-Jack, S. (2018) ‘Prediabetes and Cardiovascular Disease: Pathophysiology and Interventions for Prevention and Risk Reduction.’, *Endocrinology and metabolism clinics of North America*, 47(1), pp. 33–50. Available at: <https://doi.org/10.1016/j.ecl.2017.10.001>.
- Brož, J. et al. (2020) ‘Prevalence of diabetes and prediabetes and its risk factors in adults aged 25–64 in the Czech Republic: A cross-sectional study’, *Diabetes Research and Clinical Practice*, 170, p. 108470. Available at: <https://doi.org/10.1016/j.diabres.2020.108470>.
- Budiastutik, I. et al. (2022) ‘High Prevalence of Prediabetes and Associated Risk Factors in Urban Areas of Pontianak, Indonesia: A Cross-Sectional Study’, *Journal of Obesity*, 2022, pp. 1–9. Available at: <https://doi.org/10.1155/2022/4851044>.
- Campagna, D. et al. (2019) ‘Smoking and diabetes: dangerous liaisons and confusing relationships’, *Diabetology & Metabolic Syndrome*, 11(1), p. 85. Available at: <https://doi.org/10.1186/s13098-019-0482-2>.
- CDC (2024a) *National Diabetes Statistics Report*. Available at: https://www.cdc.gov/diabetes/php/data-research/index.html#cdc_report_pub_study_section_3-prevalence-of-diagnosed-diabetes (Accessed: 6 October 2024).
- CDC (2024b) *Prediabetes – Your Chance to Prevent Type 2 Diabetes*. Available at: <https://www.cdc.gov/diabetes/prevention-type-2/prediabetes-prevent-type-2.html> (Accessed: 12 October 2024).
- CDC (2024c) *The Surprising Truth About Prediabetes*.
- Chai, Y. et al. (2022) ‘Association of body mass index with risk of prediabetes in Chinese adults: A population-based cohort study’, *Journal of Diabetes Investigation*, 13(7), pp. 1235–1244. Available at: <https://doi.org/10.1111/jdi.13783>.
- Chowdhury, A. and S, D. (2019) ‘Prevalence and risk factor of prediabetes: A cross-sectional study among young medical students in Mangalore’, *National*

- Journal of Physiology, Pharmacy and Pharmacology*, (0), p. 1. Available at: <https://doi.org/10.5455/njppp.2019.9.0206110032019>.
- Dany, F. *et al.* (2020) ‘Urban-rural distinction of potential determinants for prediabetes in Indonesian population aged ≥ 15 years: a cross-sectional analysis of Indonesian Basic Health Research 2018 among normoglycemic and prediabetic individuals’, *BMC Public Health*, 20(1), p. 1509. Available at: <https://doi.org/10.1186/s12889-020-09592-7>.
- Deschênes, S.S. *et al.* (2023) ‘Prediabetes and the risk of type 2 diabetes: Investigating the roles of depressive and anxiety symptoms in the Lifelines cohort study’, *Diabetic Medicine*, 40(7). Available at: <https://doi.org/10.1111/dme.15061>.
- Dhawan, D. and Sharma, S. (2020) ‘Abdominal Obesity, Adipokines and Non-communicable Diseases.’, *The Journal of steroid biochemistry and molecular biology*, 203, p. 105737. Available at: <https://doi.org/10.1016/j.jsbmb.2020.105737>.
- Diabetes UK (2023) *Depression is a risk factor for type 2 diabetes, our research reveals*. Available at: <https://www.diabetes.org.uk/about-us/news-and-views/depression-risk-factor-type-2-diabetes-our-research-reveals> (Accessed: 12 October 2024).
- Diaz-Santana, M. V. *et al.* (2022) ‘Persistence of Risk for Type 2 Diabetes After Gestational Diabetes Mellitus’, *Diabetes Care*, 45(4), pp. 864–870. Available at: <https://doi.org/10.2337/dc21-1430>.
- Echouffo-Tcheugui, J.B. and Selvin, E. (2021) ‘Prediabetes and What It Means: The Epidemiological Evidence’, *Annual Review of Public Health*, 42(1), pp. 59–77. Available at: <https://doi.org/10.1146/annurev-publhealth-090419-102644>.
- Elyantari, G. *et al.* (2018) ‘Total cholesterol and HDL cholestrol as risk factor of prediabetes and diabetes in palembang city’, *Jurnal Kebidanan dan Keperawatan Aisyiyah*, 14(2), pp. 118–124. Available at: <https://doi.org/10.31101/jkk.745>.
- Fang, H. *et al.* (2018) ‘How to best assess abdominal obesity.’, *Current opinion in clinical nutrition and metabolic care*, 21(5), pp. 360–365. Available at: <https://doi.org/10.1097/MCO.0000000000000485>.
- FDA (2024) *How Smoking Can Increase Risk for and Affect Diabetes*. Available at: <https://www.fda.gov/tobacco-products/health-effects-tobacco-use/how-smoking-can-increase-risk-and-affect-diabetes> (Accessed: 6 October 2024).
- Formagini, T. *et al.* (2023) ‘Prediabetes prevalence and awareness by race, ethnicity, and educational attainment among U.S. adults’, *Frontiers in Public Health*, 11. Available at: <https://doi.org/10.3389/fpubh.2023.1277657>.

- Fujiyoshi, A. *et al.* (2016) ‘Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA)’, *Preventive Medicine Reports*, 4, pp. 225–232. Available at: <https://doi.org/10.1016/j.pmedr.2016.06.013>.
- Gaita, L. *et al.* (2024) ‘Lipid Disorders Management Strategies (2024) in Prediabetic and Diabetic Patients’, *Pharmaceuticals*, 17(2), p. 219. Available at: <https://doi.org/10.3390/ph17020219>.
- Glechner, A. *et al.* (2018) ‘Effects of lifestyle changes on adults with prediabetes: A systematic review and meta-analysis’, *Primary Care Diabetes*, 12(5), pp. 393–408. Available at: <https://doi.org/10.1016/j.pcd.2018.07.003>.
- Gong, D. *et al.* (2023) ‘From normal population to prediabetes and diabetes: study of influencing factors and prediction models’, *Frontiers in Endocrinology*, 14. Available at: <https://doi.org/10.3389/fendo.2023.1225696>.
- Gong, R. *et al.* (2021) ‘Associations between TG/HDL ratio and insulin resistance in the US population: a cross-sectional study’, *Endocrine Connections*, 10(11), pp. 1502–1512. Available at: <https://doi.org/10.1530/EC-21-0414>.
- Grundlingh, N. *et al.* (2022) ‘Assessment of prevalence and risk factors of diabetes and pre-diabetes in South Africa’, *Journal of Health, Population and Nutrition*, 41(1), p. 7. Available at: <https://doi.org/10.1186/s41043-022-00281-2>.
- Gunawan, B. *et al.* (2024) ‘Risk factors of early onset type 2 diabetes mellitus in young adults ≤ 40 years old’, *Universa Medicina*, 43(2), pp. 179–187. Available at: <https://doi.org/10.18051/UnivMed.2024.v43.179-187>.
- Hamoudi, R. *et al.* (2019) ‘Prediabetes and diabetes prevalence and risk factors comparison between ethnic groups in the United Arab Emirates’, *Scientific Reports*, 9(1), p. 17437. Available at: <https://doi.org/10.1038/s41598-019-53505-7>.
- Han, Y. *et al.* (2023) ‘Association between body mass index and reversion to normoglycemia from impaired fasting glucose among Chinese adults: a 5-year cohort study.’, *Frontiers in endocrinology*, 14, p. 1111791. Available at: <https://doi.org/10.3389/fendo.2023.1111791>.
- Howard, A.A., Arnsten, J.H. and Gourevitch, M.N. (2004) ‘Effect of Alcohol Consumption on Diabetes Mellitus’, *Annals of Internal Medicine*, 140(3), p. 211. Available at: <https://doi.org/10.7326/0003-4819-140-6-200403160-00011>.

- Hsu, J.-C. *et al.* (2023) ‘Prediabetes increases the risk of major limb and cardiovascular events’, *Cardiovascular Diabetology*, 22(1), p. 348. Available at: <https://doi.org/10.1186/s12933-023-02085-y>.
- Huang, Y. *et al.* (2016) ‘Association between prediabetes and risk of cardiovascular disease and all cause mortality: systematic review and meta-analysis.’, *BMJ (Clinical research ed.)*, 355, p. i5953. Available at: <https://doi.org/10.1136/bmj.i5953>.
- IDF (2021) *IDF Diabetes Atlas 10th edition*. Available at: https://diabetesatlas.org/idfawp/resource-files/2021/07/IDF_Atlas_10th_Edition_2021.pdf (Accessed: 6 October 2024).
- Jadhav, R.A. *et al.* (2022) ‘Measurement of visceral fat for early prediction of prediabetes-Cross-sectional study from Southern India.’, *Journal of Taibah University Medical Sciences*, 17(6), pp. 983–990. Available at: <https://doi.org/10.1016/j.jtumed.2022.05.006>.
- Jeon, E. *et al.* (2019) ‘833-P: Relationship between Prediabetes and Depression in Korean Adults’, *Diabetes*, 68(Supplement_1). Available at: <https://doi.org/10.2337/db19-833-P>.
- Karlsson, T. *et al.* (2019) ‘Contribution of genetics to visceral adiposity and its relation to cardiovascular and metabolic disease.’, *Nature medicine*, 25(9), pp. 1390–1395. Available at: <https://doi.org/10.1038/s41591-019-0563-7>.
- Kautzky-Willer, A., Harreiter, J. and Pacini, G. (2016) ‘Sex and Gender Differences in Risk, Pathophysiology and Complications of Type 2 Diabetes Mellitus.’, *Endocrine reviews*, 37(3), pp. 278–316. Available at: <https://doi.org/10.1210/er.2015-1137>.
- Kautzky-Willer, A., Leutner, M. and Harreiter, J. (2023) ‘Sex differences in type 2 diabetes’, *Diabetologia*, 66(6), pp. 986–1002. Available at: <https://doi.org/10.1007/s00125-023-05891-x>.
- Kemenkes RI (2024a) *Depresi*. Available at: <https://ayosehat.kemkes.go.id/topik-penyakit/kelainan-mental/depresi> (Accessed: 13 January 2025).
- Kemenkes RI (2024b) *Dewasa 19-59 Tahun*. Available at: <https://ayosehat.kemkes.go.id/kategori-usia/dewasa> (Accessed: 12 October 2024).
- Kemenkes RI (2024c) *Mengenal Penyakit Hipertensi*. Available at: <https://upk.kemkes.go.id/new/mengenal-penyakit-hipertensi> (Accessed: 6 October 2024).

- Kim, M.-J. *et al.* (2015) ‘Hypertension is an independent risk factor for type 2 diabetes: the Korean genome and epidemiology study.’, *Hypertension research : official journal of the Japanese Society of Hypertension*, 38(11), pp. 783–9. Available at: <https://doi.org/10.1038/hr.2015.72>.
- Kojska, I., Chacińska, M. and Błachnio-Zabielska, A. (2020) ‘Obesity, Bioactive Lipids, and Adipose Tissue Inflammation in Insulin Resistance.’, *Nutrients*, 12(5). Available at: <https://doi.org/10.3390/nu12051305>.
- Krysiak, R. *et al.* (2018) ‘Sexual functioning and depressive symptoms in women with various types of prediabetes — a pilot study’, *Endokrynologia Polska*, 69(2), pp. 175–181. Available at: <https://doi.org/10.5603/EP.2018.0021>.
- Kuang, M. *et al.* (2022) ‘Association of LDL:HDL ratio with prediabetes risk: a longitudinal observational study based on Chinese adults’, *Lipids in Health and Disease*, 21(1), p. 44. Available at: <https://doi.org/10.1186/s12944-022-01655-5>.
- Kulkarni, A., Jain, M. and Bhalerao, S. (2024) ‘General and central obesity in adult pre-diabetics: A metabolic derangement to consider’, *National Journal of Physiology, Pharmacy and Pharmacology*, 14(1), p. 1. Available at: <https://doi.org/10.5455/njppp.2023.13.11545202321122023>.
- Kusuma, I.S. and Syarif, S. (2023) ‘Predictors of Prediabetes and Undiagnosed Diabetes Mellitus Based on HbA1c Levels in Indonesia’, *Jurnal Ilmu Kesehatan Masyarakat*, 14(2), pp. 185–197. Available at: <https://doi.org/10.26553/jikm.2023.14.2.185-197>.
- Lascar, N. *et al.* (2018) ‘Type 2 diabetes in adolescents and young adults’, *The Lancet Diabetes & Endocrinology*, 6(1), pp. 69–80. Available at: [https://doi.org/10.1016/S2213-8587\(17\)30186-9](https://doi.org/10.1016/S2213-8587(17)30186-9).
- Li, Yongze *et al.* (2020) ‘Prevalence of diabetes recorded in mainland China using 2018 diagnostic criteria from the American Diabetes Association: national cross sectional study’, *BMJ*, p. m997. Available at: <https://doi.org/10.1136/bmj.m997>.
- Liao, Y.-S. *et al.* (2023) ‘Educational attainment affects the diagnostic time in type 2 diabetes mellitus and the mortality risk of those enrolled in the diabetes pay-for-performance program’, *Health Policy*, 138, p. 104917. Available at: <https://doi.org/10.1016/j.healthpol.2023.104917>.
- Liberty, I.A. and Kodim, N. (2017) ‘Assess Prediabetes Risk, As a Golden Period for Prevention of Diabetes’, *Asian Journal of Pharmaceutical and Clinical Research*, 10(6), p. 349. Available at: <https://doi.org/10.22159/ajpcr.2017.v10i6.18215>.

- Lim, J., Lee, J.A. and Cho, H.-J. (2018) ‘Association of Alcohol Drinking Patterns With Presence of Impaired Fasting Glucose and Diabetes Mellitus Among South Korean Adults.’, *Journal of Epidemiology*, 28(3), pp. 117–124. Available at: <https://doi.org/10.2188/jea.JE20170021>.
- Lizzo, J.M., Goyal, A. and Gupta, V. (2025) *Adult Diabetic Ketoacidosis*.
- Lyu, Y.S. *et al.* (2019) ‘Prevalence and Risk Factors for Undiagnosed Glucose Intolerance Status in Apparently Healthy Young Adults Aged <40 Years: The Korean National Health and Nutrition Examination Survey 2014–2017’, *International Journal of Environmental Research and Public Health*, 16(13), p. 2393. Available at: <https://doi.org/10.3390/ijerph16132393>.
- Maghfiroh, A.A., Simanjorang, C. and Karima, U.Q. (2024) ‘Factors Associated with the Incidence of Prediabetes in Bogor, Indonesia: A Cohort Study’, *Journal of Research in Health Sciences*, 25, p. e00635. Available at: <https://doi.org/10.34172/jrhs.2025.170>.
- Masri, E., Harleni and Zebua, I. (2024) ‘Gambaran Pengetahuan dan Konsumsi Makanan Berisiko pada Mahasiswa S1 Gizi Universitas Perintis Indonesia Tahun 2023’, *Ensiklopedia of Journal*, 6(4), pp. 35–46.
- Mehdi, S. *et al.* (2023) ‘A review on linking stress, depression, and insulin resistance via low-grade chronic inflammation’, *Biochemistry and Biophysics Reports*, 36, p. 101571. Available at: <https://doi.org/10.1016/j.bbrep.2023.101571>.
- Menteri Kesehatan (2013) *Permenkes No. 30 Tahun 2013*.
- Menteri Kesehatan RI (2016) *PERMENKES RI No 25 Tahun 2016*.
- Moran, L.J. *et al.* (2010) ‘Impaired glucose tolerance, type 2 diabetes and metabolic syndrome in polycystic ovary syndrome: a systematic review and meta-analysis’, *Human Reproduction Update*, 16(4), pp. 347–363. Available at: <https://doi.org/10.1093/humupd/dmq001>.
- Mourouti, N. *et al.* (2023) ‘The Association of Lifestyle Patterns with Prediabetes in Adults from Families at High Risk for Type 2 Diabetes in Europe: The Feel4Diabetes Study’, *Nutrients*, 15(14), p. 3155. Available at: <https://doi.org/10.3390/nu15143155>.
- Mujiono, M., Udijono, A. and Kusuma, D. (2023) ‘Faktor-Faktor yang Berhubungan dengan Kejadian Prediabetes’, *MEDIA KESEHATAN MASYARAKAT INDONESIA*, 22(5), pp. 314–318. Available at: <https://doi.org/10.14710/mkmi.22.5.314-318>.
- Nasution, H., Febriyanti, E. and Suryani, D. (2022) ‘Relationship between Frequency of Sugar Sweetened-Beverages (SSB) Consumption and

- Prediabetes: Aim For Screening Prediabetes Among Medical Students', *Buletin Farmatera*, 7(1), pp. 51–57. Available at: <https://doi.org/https://doi.org/10.30596/bf.v7i1.9439.g8814>.
- NCEP ATP III (2003) 'The National Cholesterol Education Program Adult Treatment Panel III Guidelines', *Journal of Managed Care Pharmacy*, 9(1 Supp A), pp. 2–5. Available at: <https://doi.org/10.18553/jmcp.2003.9.s1.2>.
- Nordström*, A. et al. (2016) 'Higher Prevalence of Type 2 Diabetes in Men Than in Women Is Associated With Differences in Visceral Fat Mass', *The Journal of Clinical Endocrinology & Metabolism*, 101(10), pp. 3740–3746. Available at: <https://doi.org/10.1210/jc.2016-1915>.
- Olivine, A. (2024) *Symptoms of Prediabetes in Females*. Available at: <https://www.verywellhealth.com/symptoms-of-prediabetes-in-females-6891606> (Accessed: 6 October 2024).
- Park, J.C. et al. (2023) 'Association of Sustained Low or High Income and Income Changes With Risk of Incident Type 2 Diabetes Among Individuals Aged 30 to 64 Years', *JAMA Network Open*, 6(8), p. e2330024. Available at: <https://doi.org/10.1001/jamanetworkopen.2023.30024>.
- Park, K.S. and Hwang, S.Y. (2020) 'Lifestyle-related predictors affecting prediabetes and diabetes in 20-30-year-old young Korean adults.', *Epidemiology and health*, 42, p. e2020014. Available at: <https://doi.org/10.4178/epih.e2020014>.
- Pazmino, L. et al. (2021) 'Impact of Work and Recreational Physical Activity on Prediabetes Condition among U.S. Adults: NHANES 2015–2016', *International Journal of Environmental Research and Public Health*, 18(4), p. 1378. Available at: <https://doi.org/10.3390/ijerph18041378>.
- Peng, J. et al. (2021) 'Association between dyslipidemia and risk of type 2 diabetes mellitus in middle-aged and older Chinese adults: a secondary analysis of a nationwide cohort.', *BMJ open*, 11(5), p. e042821. Available at: <https://doi.org/10.1136/bmjopen-2020-042821>.
- Pranata, S. and Khasanah, D. (2017) *Merawat Penderita Diabetes Melitus*. Yogyakarta: Pustaka Panasea.
- Pratiwi, I.N. et al. (2024) 'Predictors of Prediabetes Among Young Adults in East Java, Indonesia: A Cross-sectional Study', *Nurse Media Journal of Nursing*, 14(2). Available at: <https://doi.org/10.14710/nmjn.v14i2.56904>.
- Puustinen, P.J. et al. (2011) 'Psychological Distress Predicts the Development of the Metabolic Syndrome: A Prospective Population-Based Study', *Psychosomatic Medicine*, 73(2), pp. 158–165. Available at: <https://doi.org/10.1097/PSY.0b013e3182037315>.

- Qi, Y. *et al.* (2019) ‘Adulthood Socioeconomic Position and Type 2 Diabetes Mellitus-A Comparison of Education, Occupation, Income, and Material Deprivation: The Maastricht Study.’, *International journal of environmental research and public health*, 16(8). Available at: <https://doi.org/10.3390/ijerph16081435>.
- Salas-Salvadó, J. *et al.* (2019) ‘Effect of a Lifestyle Intervention Program With Energy-Restricted Mediterranean Diet and Exercise on Weight Loss and Cardiovascular Risk Factors: One-Year Results of the PREDIMED-Plus Trial’, *Diabetes Care*, 42(5), pp. 777–788. Available at: <https://doi.org/10.2337/dc18-0836>.
- Sallar, A. and Dagogo-Jack, S. (2020) ‘Regression from prediabetes to normal glucose regulation: State of the science’, *Experimental Biology and Medicine*, 245(10), pp. 889–896. Available at: <https://doi.org/10.1177/1535370220915644>.
- Schmidt, C. *et al.* (2023) ‘Psychosocial work conditions and prediabetes risks: a cross-sectional study in middle-aged men and women’, *Scientific Reports*, 13(1), p. 1198. Available at: <https://doi.org/10.1038/s41598-023-28420-7>.
- Schweitzer, G.G. *et al.* (2023) ‘Elevated liver glycogenolysis mediates higher blood glucose during acute exercise in Barth syndrome’, *PLOS ONE*, 18(8), p. e0290832. Available at: <https://doi.org/10.1371/journal.pone.0290832>.
- Shen, Y. *et al.* (2018) ‘Gestational diabetes with diabetes and prediabetes risks: a large observational study’, *European Journal of Endocrinology*, 179(1), pp. 51–58. Available at: <https://doi.org/10.1530/EJE-18-0130>.
- Shou, J., Chen, P.-J. and Xiao, W.-H. (2020) ‘Mechanism of increased risk of insulin resistance in aging skeletal muscle.’, *Diabetology & metabolic syndrome*, 12, p. 14. Available at: <https://doi.org/10.1186/s13098-020-0523-x>.
- Soelistijo, S. *et al.* (2021) *Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 Dewasa di Indonesia - 2021*. PB Perkeni.
- Sudibjo, P., Rismayanthi, C. and Apriyanto, K.D. (2021) ‘Hubungan antara sindrom metabolik dengan kebugaran jasmani pada lansia’, *Jurnal Keolahragaan*, 9(2), pp. 159–167. Available at: <https://doi.org/10.21831/jk.v9i2.41007>.
- Sukenty, N.T., Shaluhiyah, Z. and Suryoputro, A. (2018) ‘Faktor Perilaku dan Gaya Hidup yang Mempengaruhi Status Prediabetes Pasien Puskesmas Pati II’, *Jurnal Promosi Kesehatan Indonesia*, 13(2), p. 129. Available at: <https://doi.org/10.14710/jPKI.13.2.129-142>.
- Sun, J. *et al.* (2022) ‘The Correlation of Prediabetes and Type 2 Diabetes With Adiposity in Adults’, *Frontiers in Nutrition*, 9. Available at: <https://doi.org/10.3389/fnut.2022.818263>.

- Syauqy, A., Mattarahmawati, S.A. and Pramono, A. (2022) ‘Food Consumption in Relation to Hyperglycemia in Middle-Aged Adults (45–59 years): A Cross-Sectional National Data Analysis’, *Jurnal Gizi dan Pangan*, 17(3), pp. 187–194. Available at: <https://doi.org/10.25182/jgp.2022.17.3.187-194>.
- Tabák, A.G. *et al.* (2012) ‘Prediabetes: a high-risk state for diabetes development.’, *Lancet (London, England)*, 379(9833), pp. 2279–90. Available at: [https://doi.org/10.1016/S0140-6736\(12\)60283-9](https://doi.org/10.1016/S0140-6736(12)60283-9).
- Tangjittipokin, W. *et al.* (2022) ‘Prevalence and Characteristics of Prediabetes and Metabolic Syndrome in Seemingly Healthy Persons at a Health Check-Up Clinic’, *Journal of Multidisciplinary Healthcare*, Volume 15, pp. 1585–1594. Available at: <https://doi.org/10.2147/JMDH.S374164>.
- Tian, Y. *et al.* (2024) ‘Associations of Diabetes and Prediabetes With Mortality and Life Expectancy in China: A National Study’, *Diabetes Care*, 47(11), pp. 1969–1977. Available at: <https://doi.org/10.2337/dca24-0012>.
- Tramunt, B. *et al.* (2020) ‘Sex differences in metabolic regulation and diabetes susceptibility’, *Diabetologia*, 63(3), pp. 453–461. Available at: <https://doi.org/10.1007/s00125-019-05040-3>.
- Tsimihodimos, V. *et al.* (2018) ‘Hypertension and Diabetes Mellitus’, *Hypertension*, 71(3), pp. 422–428. Available at: <https://doi.org/10.1161/HYPERTENSIONAHA.117.10546>.
- Urena-Bogarin, E.L. *et al.* (2015) ‘Prevalence of pre-diabetes in young Mexican adults in primary health care’, *Family Practice*, 32(2), pp. 159–164. Available at: <https://doi.org/10.1093/fampra/cmu047>.
- Volpe, M. *et al.* (2023) ‘The Emerging Role of Prediabetes and Its Management: Focus on L-Arginine and a Survey in Clinical Practice.’, *High blood pressure & cardiovascular prevention : the official journal of the Italian Society of Hypertension*, 30(6), pp. 489–496. Available at: <https://doi.org/10.1007/s40292-023-00613-1>.
- Wagner, R. *et al.* (2013) ‘Family history of diabetes is associated with higher risk for prediabetes: a multicentre analysis from the German Center for Diabetes Research’, *Diabetologia*, 56(10), pp. 2176–2180. Available at: <https://doi.org/10.1007/s00125-013-3002-1>.
- Wang, D. *et al.* (2022) ‘Association between visceral adiposity index and risk of prediabetes: A meta-analysis of observational studies.’, *Journal of diabetes investigation*, 13(3), pp. 543–551. Available at: <https://doi.org/10.1111/jdi.13685>.
- Weir, C.B. and Jan, A. (2024) *BMI Classification Percentile And Cut Off Points*.

- WHO (2006) *Definition and Diagnosis Of Diabetes Mellitus and Intermediate Hyperglycemia, WHO Library Cataloguing-in-Publication Data*. Available at: https://iris.who.int/bitstream/handle/10665/43588/9241594934_eng.pdf (Accessed: 6 October 2024).
- WHO (2021) *Global Physical Activity Questionnaire (GPAQ) Analysis Guide*. Available at: <https://www.who.int/docs/default-source/ncds/ncd-surveillance/gpaq-analysis-guide.pdf> (Accessed: 16 October 2024).
- Wu, Y. *et al.* (2015) ‘Activation of AMPK α 2 in adipocytes is essential for nicotine-induced insulin resistance in vivo’, *Nature Medicine*, 21(4), pp. 373–382. Available at: <https://doi.org/10.1038/nm.3826>.
- Xie, J. *et al.* (2022) ‘Global burden of type 2 diabetes in adolescents and young adults, 1990–2019: systematic analysis of the Global Burden of Disease Study 2019’, *BMJ*, p. e072385. Available at: <https://doi.org/10.1136/bmj-2022-072385>.
- Yang, T. *et al.* (2022) ‘Correlation between the triglyceride-to-high-density lipoprotein cholesterol ratio and other unconventional lipid parameters with the risk of prediabetes and Type 2 diabetes in patients with coronary heart disease: a RCSCD-TCM study in China.’, *Cardiovascular diabetology*, 21(1), p. 93. Available at: <https://doi.org/10.1186/s12933-022-01531-7>.
- Yang, W. *et al.* (2024) ‘Different levels of physical activity and risk of developing type 2 diabetes among adults with prediabetes: a population-based cohort study’, *Nutrition Journal*, 23(1), p. 107. Available at: <https://doi.org/10.1186/s12937-024-01013-4>.
- Yesmin, M., Ali, M. and Saha, S. (2023) ‘The prevalence and influencing factors of coexisting prediabetes and prehypertension among Bangladeshi adults’, *BMC Public Health*, 23(1), p. 1184. Available at: <https://doi.org/10.1186/s12889-023-16090-z>.
- Yokoyama, H. *et al.* (2007) ‘Effects of Excessive Ethanol Consumption on the Diagnosis of the Metabolic Syndrome Using Its Clinical Diagnostic Criteria’, *Internal Medicine*, 46(17), pp. 1345–1352. Available at: <https://doi.org/10.2169/internalmedicine.46.6196>.
- Yun, J.E. *et al.* (2012) ‘Smoking Is Associated With Abdominal Obesity, Not Overall Obesity, in Men With Type 2 Diabetes’, *Journal of Preventive Medicine & Public Health*, 45(5), pp. 316–322. Available at: <https://doi.org/10.3961/jpmph.2012.45.5.316>.
- Zand, A., Ibrahim, K. and Patham, B. (2018) ‘Prediabetes: Why Should We Care?’, *Methodist DeBakey cardiovascular journal*, 14(4), pp. 289–297. Available at: <https://doi.org/10.14797/mdcj-14-4-289>.

- Zheng, T., Gao, Y. and Tian, H. (2012) ‘Relationship between blood lipid profiles and pancreatic islet β cell function in Chinese men and women with normal glucose tolerance: a cross-sectional study’, *BMC Public Health*, 12(1), p. 634. Available at: <https://doi.org/10.1186/1471-2458-12-634>.
- Zhuang, Q. et al. (2020) ‘Negative emotions in community-dwelling adults with prediabetes and hypertension’, *Journal of International Medical Research*, 48(4). Available at: <https://doi.org/10.1177/0300060520918411>.