

DAFTAR PUSTAKA

Appley,A.G and Louis Solomon.(1995).Terjemahan Ortopedi dan Fraktur Sistem Appley (edisi ke7).Widya Medika

Australian Journal of Physiotherapy 2005 Vol. 51 “Do early quadriceps exercises affect the outcome of ACL”

Allison A. Hall, Brenda L. Schrage , 1995 .”Comparison of Isometric Quadriceps Setting Techniques: An Electromyographic Study “Pacific University.

Baker PA, Evans OM, Lee C (1991). Treadmill gait retraining following fractured neck of femur. Arch Phys Med Rehabil;72:649-52.

Smeltzer SC. 2001. Buku Ajar Keperawatan Medikal Bedah. Jakarta EGC.
Hal : 45-47.

Ceder L, Svensson K, Thorngren K-G. 1980 Statistical prediction of rehabilitation in elderly patients with hip fractures. Clin Orthop 152:185- 90.

Cummings SR, 1995. Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. N Engl J Med.;332:/767_73.

Cameron I, Crotty M, Currie C, 2000. Geriatric rehabilitation following fractures in older people: a systematic review. Health Technol Assess 4:1-111.

Dorland, Newman. 2002. *Kamus Kedokteran Dorland*. Edisi 29,
Jakarta:EGC,1765.

Don W.Fwacett. 2002. Buku ajar Histologi. Edisi 12, Jakarta, EGC.

Dreeben-Irimia , 2011 ."Introduction to Physical Therapy for Physical Therapist Assistants".

Evelyn C, Pearce. (2011). *Anatomi dan Fisiologi untuk Paramedis*. Jakarta: PT Gramedia Pustaka Utama.

Gross,Jeffery M .2009.*muskuloskeletal examination*. USA

Higgins, Michael, 2011 "Therapeutic Exercise: From Theory to Practice" Hal- 449 Quadricep setting.

Jones and barlett, 2003. Anatomi fisiologi untuk pemula, Jakarta : EGC.

Knee Strength. Journal Of Estide Sport, Rehabilatasi Clinic (*unduh mei 2014*)

Kannus P, Niemi. 1970. Hip fractures in Finland between 1970 and 1997
and predictions for the future Lancet. 353:6:/802_5

Lamb SE, 1995. Mobility after proximal femoral fracture: the relevance of leg extensor power, postural sway and other factors. Age Ageing 24:308-14.

Lippincott Williams & Wilkins (2006), Rockwood&Green Patah tulang pada Dewasa ,*edisi 6, 47*

Mitchell SL, 2001. Randomized controlled trial of quadriceps training after proximal femoral fracture. Clin Rehabil 15:282-90.

Nguyen ND,2005. Identification of high-risk individuals for hip fracture: A 14-year prospective study. J Bone Miner Res, 20:/1921_8.

Nevalainen, 2004. Functional ability after hip fracture among home-dwelling at the time a fracture.Cent Eur Public Health 12:/211_6.

Osnes EK, Lofthus CM,Meyer HE, 2004. Falch JA, Nordsletten L, Cappelen I, et al Consequences of hip fracture on activities of daily life and residential needs. Osteoporos Int. ; 15: 567_74. Van Balen R, Essink-Bot ML, Steyberg E, Cools H, Habbema DF. Quality of life after hip fracture: A comparison of four.

Parker MJ, Handoll HH, Dynan Y,2000. Mobilisation strategies after hip fracture surgery in adults. Cochrane Database Syst Rev CD001704.

R.Puts & R.Pabs (2003), Sobotta, Edisi 21, 276-311

Reeves CJ, Roux G and Lockhart R, 2001, Keperawatan Medikal Bedah, Buku I, (Penerjemah Joko Setyono), Jakarta : Salemba Medika

Rosell PA, Parker MJ.2003.Functional outcome after hip fracture. A 1-year prospective outcome study of 275 patients. Injury 34:7:/529_32.

Repatriation General Hospital, Daw Park, Australia.2002. "Early discharge and home rehabilitation after hip fracture achieves functional improvements: a randomized controlled trial "*Clinical Rehabilitation*; 16: 406–413

Shepperd S, Iliffe S.2000. Hospital-at-home versus in-patient hospital care. Cochrane Database Syst Rev 2:CD000356.

Stevens JA, Olson S. 2000. Reducing falls and resulting hip fractures among older women. MMWR Recomm Rep. 31:/3_12.

Sherrington C, Lord S.2005. Reliability of simple portable tests of physical performance in older people after hip fracture.
Clin Rehabil.; 19: 496_504

The journal Houghton Company Mifflin. (2004 - 2007), The American Heritage medical Dictionary

Tinetti ME, Baker DI, Gottschalk M,1987. Systematic home-based physical and functional therapy for older persons after hip fracture.
Arch Phys Med Rehabil 78:1237-47.

The journal 2008.The effects of muscle strength and power training on mobility among older hip fracture patients". Advances in Physiotherapy.10:195_202

Tinetti ME, Baker DI, Gottschalk M, 1997. Systematic home-based physical and functional therapy for older persons after hip fracture
Arch Phys Med Rehabil78:1237-47.

2014 ; Gambar Tipe Fraktur (*Doctorology.net,unduh mei 2014*)

2014 : Gambar bentuk latihan quadriceps (www.mmg.com unduh mei 2014)